STATE OF ALASKA

To: Distribution Date: 9-16-87

Disk:C:\Word\FY88
File:Index.sep

From: Steve Elliott Telephone: 465-4270

Fishery Biologist Sport Fish Division

Region I

Leon Shaul

Fishery Biologist

Commercial Fisheries Division

Region I

SUBJECT: COHO SALMON SPAWNER INDEX STREAMS

Attached is the finalized list of coho index streams for 1987 based on your comments and recommendations. Between the Sport Fish Division, U.S. Canada, and Commercial Fisheries Divison budgets we should have adequate funding for all surveys. Please make a final review; call Leon or myself if you have any changes.

Attachment

Distribution: Greg Thomason, CF, Yakutat

Ray Staska, CF, Haines
Don Ingledue, CF, Juneau
Doug Jones, CF, Juneau
Bob DeJong, CF, Sitka

Will Bergman, CF, Petersburg Robert Larson, CF, Petersburg Steve Hoffman, SF, Ketchikan Dennis Hubartt, SF, Ketchikan

Art Schmidt, SF, Sitka Mike Bethers, SF, Juneau Randy Erikson, SF, Haines Bob Johnson, SF, Yakutat Bob Zorich, SF, Petersburg

cc: Dave Cantillon
Gary Gunstrom
Gary Sanders

Date last revised:9-15-87

Coho Salmon Escapement Index Streams to be surveyed September - November, 1987 and agency responsible for survey. (1 = hatchery influence; 2 = possible straying from hatchery releases; 3=Survey by Commercial Fisheries Div staff -- funding by Sport Fish Div.)

HAINES AREA:

| Chilkat River (20-22 mi) | 115-32-025 | (SF) |
|--------------------------|------------|------|
| Clear Creek | 115-32-027 | (SF) |
| Takhin River | 115-32-030 | (CF) |
| Spring Creek | 115-32-040 | (SF) |
| 31 mile Creek | 115-32-057 | (SF) |
| Kelsall River | 115-32-064 | (SF) |
| Tahini River | 115-32-068 | (SF) |

JUNEAU AREA:

| Switzer Creek | 111-40-007 | (SF) |
|-------------------|------------|-------|
| Peterson Creek | 111-50-010 | (SF) |
| Auke Creek (weir) | 111-50-042 | (ABL) |
| Montana Creek 2 | 111-50-052 | (SF) |
| Steep Creek | 111-50-056 | (SF) |
| Jordan Creek | 111-50-062 | (SF) |
| Outer Point Creek | 111-50-075 | (SF) |
| Hasselborg River | 112-67-035 | (CF) |
| Chiak Bay Creek | 112-80-028 | (CF) |
| Berners River | 115-20-010 | (CF) |

Taku River Tributaries:

| Sockeye Creek | 111-32-038 | (SF&CF) |
|----------------------|------------|---------|
| Fish Creek | 111-32-056 | (SF&CF) |
| Flannagan Slough | 111-32-048 | (SF) |
| Yehring Creek (weir) | 111-32-066 | (SF&CF) |
| Johnson Creek | 111-32-068 | (SF&CF) |
| Nahlin River | 111-32-270 | (CF) |
| Dudidontu River | 111-32-280 | (CF) |

KETCHIKAN AREA:

| Tombstone River | 101-15-019 | (CF) |
|-------------------------|------------|---------|
| Fish Creek | 101-15-085 | (CF) |
| Humpback Creek | 101-30-083 | (CF) |
| White River | 101-45-024 | (CF&SF) |
| Carrol River | 100-45-078 | (SF) |
| Ward Creek ¹ | 101-47-015 | (SF) |
| Indian River | 101-71-004 | (SF) |
| Walker Creek | 101-71-028 | (CF) |
| Eulachon River | 101-75-015 | (SF) |

| Herman Creek Traitors Creek | 101-75-005 101-90-029 | (CF) (CF) |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|
| Prince of Wales Island: Twelvemile Creek Harris River Maybeso Creek Natzuhini Creek St. Nicholas Creek Cable Creek S. Staney Creek 108 Creek | 102-60-072 102-60-082 102-60-084 103-40-035 103-60-059 103-60-077 103-90-042 106-30-080 | (SF) (CF) (SF) (SF) (SF) (SF) (SF) |
| PETERSBURG AREA: | | |
| Falls Creek Bear Creek Duncan Creek Mosman Creek Mosman Creek Navy Creek Snake Creek (opt) Martin Creek (opt) Harding River N. Bradfield River E. Bradfield River Sumner Creek Ohmer Creek Kake Bake Creek Unnammed Creek (opt) Unnammed Creek (opt) Unnammed Creek (opt) Portage Creek (opt) | 106-44-006 108-50-003 106-43-075 106-22-066 106-22-016 107-30-70 107-40-38 107-40-49 107-40-52 107-40-53 108-40-40 108-40-50 109-32-020 109-42-40 109-45-10 109-52-08 110-16-002 | (SF&CF) (SF&CF) (CF) (CF) (CF) (CF) (CF) (CF) (CF/SF) (CF/SF) (CF) (CF) (CF) (CF) (CF) (CF) (CF) |
| Stikine River Tribs: North Arm Creek Shakes Slough Ketili Creek Kikahe River Goat Creek Shuktusa Creek Andrews Creek Katete Creek | 108-40-010 108-40-013 108-40-014 108-40-016 108-40-017 108-40-018 108-40-020 108-70-009 | (CF) (CF) (CF) (CF) (CF) (CF) (CF) |
| SITKA AREA: | | |
| Starragavan Creek Indian River ¹ Salmon Lake Creek Kizuchia Creek Nakwasina River Sinitsin Creek ² | 113-41-015 113-41-019 113-41-032 113-41-042 113-43-002 113-62-008 | (SF) (SF) (SF) (SF) (SF) |

••

| St. Johns Creek Black River | 113-66-006 113-81-011 | (SF) (SF) |
|--------------------------------|--------------------------|--------------|
| YAKUTAT AREA: | | |
| Airport ditches | NA | (SF) |
| Akwe River | 182-40-010 | (SF) |
| Italio River | 182-50-010 | (SF) |
| Old Situk River | 182-70-015 | (SF) |
| Situk River | 182-80-010 | (SF) |
| Tawah/Lost River | 182-80-030 | (SF) |
| Yahtse River | 185-10-010 | (CF) |
| Tsiu River | 192-42-020 | (CF) |
| Tsivat River | 192-42-040 | (CF) |
| Kaliakh River | 192-41-010 | (CF) |

MEMORANDUM STATE OF ALASKA

To: Gary Sanders Date: 7-27-87

Regional Research Supervisor

Sport Fish Div., Reg. I Disk:D:\

File:Johnson.jul

From: Steve Elliott

Fishery Biologist

Sport Fish Div., Reg. I Telephone: 465-4270

SUBJECT: YAKUTAT COHO INDEX STREAMS

Bob Johnson asked me to relay the following to you in regard to your recent memo on coho index streams. He recommends the following streams and Divisional resposibilities for the Yakutat area:

| Airport ditches | (SF) | NA |
|--------------------|------|------------------|
| Old Situk River | (SF) | 182-70-015 |
| Situk River | (SF) | 182-80-010 |
| Tawah/Lost River | (SF) | 182-80-030 |
| Akwe River | (CF) | 182-40-010 |
| Italio River | (CF) | 182-50-010 |
| Tsiu/Tsivat Rivers | (CF) | 192-42-020 & 040 |

Greg Thomason may want to add additional streams.

cc B. Johnson, SF

G. Thomason, CF

M. Bethers, SF

Steve: Kaliakh River (CF) _ 192-41 Yaktse River (CF) _ 185-10

They Homosor

State of Alaska

TO: Distribution

DATE:

February 24, 1987

FILE NO .:

THRU:

TELEPHONE NO.:

465-4270

SUBJECT:

Coho Escapement Index

Streams

FROM:

Steve Elliott Fishery Biologist
Sport Fish Division
Douglas

w/ Original Mile Bothers

The following is excerpted from the draft Sport Fish Coho Research Operation Plan. The plan is a take-off on the Coop. CF. SF. Coho Research Plan and seeks to establish a standard set of coho index streams that are surveyed each year using the same methods. The goal is consistency. Please review the plan text with the following in mind:

- Are these streams realistic index streams? Can they be done each year without fail, or are they surveys that you do only if you have the funds, time, etc (we never have good weather).
- 2. If each stream is realistic, what portion of the stream do you survey. Please be specific; where do you begin the survey and where do you end; how many miles are involved? Listing "entire" is not enough.
- 3. What is the best time for doing the survey. Notes on conditions would also be helpful. For example in the Taku, we have to wait until a good cold snap has reduced turbidity but if one waits too long the fish are scattered and impossible to count. Describing the window for optimum conditions can be important.

The index streams, area of survey, and preferred time of survey are listed on Table 1.

Al Bingham will soon be breathing down my neck for this coho operational plan. I'd appreciate your prompt attention to this and return your comments to me by March 9, 1987.

Attachment

cc: M.Bethers

R.Ericksen

B.Johnson

A.Schmidt

G.Sanders

D.Siedelman

Table 1. Location, schedule, and date of spawner escapement indiced in coho salmon index streams.

he hall my b.
Harden seb.
Hard

| tream | Survey Area | Date | Method |
|----------------------------------------------|--------------------|------------|------------|
| Maines Area: | | | |
| l mile Creek | entire | Oct-Nov | foot, |
| 7 mile Creek | entire | Oct-Nov | foot |
| Bear Creek | entire | Oct-Nov | foot |
| Chilkat River | 20-22 mile | Oct-Nov | foot,air |
| Chilkat River | (above Tahini) | Oct-Nov | foot,boat |
| lerman Creek | entire | Oct-Nov | foot |
| Kelsall River | 2 mi below bridge | Oct-Nov | foot |
| ittle Salmon R. | entire | Oct-Nov | foot |
| Nataga River | l mi. above bridge | Oct-Nov | foot |
| Spring Creek | entire | Oct-Nov | foot |
| Tahini River | from mouth | Oct-Nov | foot, boat |
| | to border. | | |
| Juneau Area: | | | |
| Auke Creek (weir) | | | |
| Jordan Creek | | | |
| Iontana Creek | | | |
| Outer Point Creek | | | |
| Peterson Creek | | | |
| Steep Creek | | | |
| Switzer Creek | | | |
| Ketchikan Area: | | | |
| Carrol River | | | |
| Eulachon River | | | |
| Indian River | | | |
| lard Creek | | | |
| Petersburg Area: | | | |
| Bear Creek | | | |
| alls Creek | | | |
| hmer Creek | | | |
| etersburg Creek | | | |
| Sumner Creek | | | |
| Sitka Area: | | | |
| Indian River | | | |
| indian kiver Kizuchia Creek | | | |
| dalana a da a Didana a | | | |
| akwasina kiver Salmon Lake Creek | | | |
| Sinitsen Creek | | | 1 184- |
| tarrigavan Creek | aul Boh Jeho | . (ALUICA | |
| starrigavan Creek St. Joh <u>ns Creek</u> | - Bab Jaho | me () | |
| akutat Area: | all I | | |
| akutat Area: | | 10/1-2/1 | foot |
| Airport dittnes Ankau River | | 8/5-10/3 | |
| Akwe River | | 9/15-10/3 | |
| italio River | | 9/15-10/ | |
| LALIO KIVEI | | J/ LJ - C/ | - 4 |

-Continued-

Table 1. (cont'd) Location, schedule, and date of spawner escapement indiced in coho salmon index streams.

| Stream | Survey Area | Date | Method |
|------------------------------------|------------------------------------------------------|-------------------------|---------------|
| Old Situk River | 10 mi. Bridge to Situk then up to 9 mi. bridge | 10/10-10/31 | float |
| Ophir Creek |) mr. 011660 | 10/1-11/15 | foot float |
| Situk River Tawah Creek/Lost R. | | 8/7-10/15 9/15-10/31 | air |
| Tsiu River | | 9/15-10/31 | air |

Coho Inchy Sheams

Stare

This is in response to your inquiry of 2/24/87 on colo index strong in the

- 1. Hes the obsame me realistic representatives of our local sheam types and they can be done on an overreal basis.
- 2. Pull Cuch weri-count all the mothers!

Jordan Creek. Two pers. one valuela. One new drops of the pers of at Gandikin Dince and then drives up to nance, sheet and paths can and counts apsheam to headwaters. Most downsheam person counts from Gandukin Drive upsheam to nance, sheet and then picks up can and drives up to Thurder Mit havier court to pick up other person. The entire sheam upsheam Kom Gandukin Drive is surveyed. See Map.—

Montana Cruk. Two pairs people, Two vehicles, to purvey in one day. Our vehicle dupped off at Back loop Bridge. Four pers in other can drive to old broken bridge and let out 2 surveyers and drive can back down to rifle range bridge

One pair people court repshram. From brother by the Up sheam to Montana Cr. trail bute. Then come back down and go up McComes Crech (incl smy pored creck) until you run out of fish. Then come track down M'Ginnis Creck and cut over to headwaters ? "Little Mc Cominis" and 30 down to confluence of Montana Crock. From there jo down to rithe range bridge, pick up can and go home and pick out decits chibs and have The other pair of people court from the 1. The range midge Aburnshenn to Back Loop midge - Maided area behind 3 814 hours and trib across from 5+1+ house, look in deep needs and log jams real well in lower area. water is way dark! The entre shear water from Back loop Bute 15 counted See Map -

Rome and one manistern - 12-2/3 up to end of shear

Peterson Crech , 25 mile. One pers can do it. Count uptream dom Glacier Hovy rup into gorge where 400'll run out of fish. Then head back downsheam and get other channels of braids. - anamarches. Count entre sheam upbshemm hom Glacier Hovy.

Steep Creek, Park at the salmon Lite history, sign Furnout, court up sheam to falls them go tack down to sheam mouth and count upshrow to can Count pools meal carefully: they are main holding arms. Good glasses are a must here. See kinp

Switzer Creck. Park can at 018 61. Hury cultures
Court down to Egan mine. Do a 10+ of poking
under under cut banks = good hiding places. Go
back upsham and count from 018 Glacien Hury
upsham to old losging on the and culturet on
hountain ane nice. Hile pheets back to can be hupof two pers avail, one start at hisdwaters and
work down, one start as out lined allows and
meet in the middle section.

| 5 dream | No People | Time For survey | Comment |
|----------------|-----------|-------------------|-------------------|
| Tordan | 2 | 3 A15 | Watch out for - |
| | 1 | 4-5 | |
| Montana Cr | 2 | 1 extalog day - 2 | persmin in |
| | 2 prs | 1 day | upper section |
| Outer Point Co | 1 | 1 day | |
| | 2 | 4-5 9,15 | |
| Peterson Cr | 1002 | | I he driving time |
| Steep or | / | 1-1/2 hrs | |
| Su. 34 G. | / | 4 h15 | |
| | 7 | 2-3 hrs | |

Would recommend fire arms be carried at Reterior a, prontana and Diter point Creek. Have seen terred black hears over the opens on middle action of gordan and lots of argue on on ou. 3, . Pack a big day street on gordan a fordance.

In all awas, study deep dack pools (tanic water) were carefully and also under 'of James etc.

sank where there enough come for a smeaky coho to hide.




Figure 46.1. Map of Peterson Creek.




Figure 47.1. Map of Peterson (Outer Point) Creek.




Figure 56.1. Map of Steep Creek.




Figure 58.1. Map of Switzer Creek.

58.




Figure 37.1. Map of McGinnis Creek.




Figure 28.1. Location of Jordan Creek.

One person can nawn in 15 mm.
The entire sheam he carre of
work up sheam he carre of
work and such ment is purished

State of Alaska

TO: Steve Elliott Fisheries Biologist Sport Fish Division Juneau DATE: July 18, 1987

FILE NO:

TELEPHONE NO: 766-2625

SUBJECT: Haines Coho Index Streams


W/MAPS


FROM: Randy Ericksen Fisheries Biologist
Sport Fish Division
Haines


Sixty to seventy percent of my time in the fall is committed to creel surveys. Add to this my other management duties and I do not realistically have the time to conduct escapement surveys on eleven coho systems. Many of the streams surveyed in the past did not have sufficient spawners to justify the time and money involved. I recommend the following systems be used as index streams:


- 1. Chilkat River (20-22 mile), 115-32-10250, This system should be surveyed by air late in the fall once the river clears. Normally Ray flies this to enumerate fall chums and counts coho concentrated near the Tsirku fan. We should survey this at least one additional time.
- 2. Spring Creek, 115-32-10250-2019-3002-4007-5002, I normally walk this stream in mid October. It is easy to hike and always seems to have good visibility.
- 3. Thirtyone mile Creek, 115-32-10250-2023-3010, I walk this stream in late October. The stream has several tributaries which are all spring fed so visibility is good.
- 4. Kelsall River, 115-32-10250-2045, I walk this system in late October and early November once the river clears. However the water is deep and surveying conditions are not always ideal. It would be better to survey this system by air.
- 5. Tahini River, 115-32-10250-2057, This system is normally surveyed by foot but takes a jet boat to reach the river. Occasionally the Chilkat river gets too low to use a jet boat so we need to charter an airboat. The river itself is subject to fall flooding which makes surveys impossible. The river should be surveyed in mid to late October.
- 6. Clear Creek, 115-32-10280, This system is surveyed by foot but is accessed by jet boat. The creek is well up Murphy Flats on the Chilkat river and takes a sizeable tide to get to. The creek should be surveyed in late October.


I recommend you budget for one day of airboat charter (about \$300) and 3hrs air charter (about \$700).














State of Alaska

TO: Gary Sanders
Acting Regional Supervisor
Sport Fish Division
Juneau

FILE NO:

FROM: Randy Ericksen

TELEPHONE NO: 766-2625

Fisheries Biologist Sport Fish Division SUBJECT: Coho Index Streams

DATE: July 28, 1987

Haines

In reviewing your proposed coho index streams for the Haines area, I noted several systems which had very few spawners, or were close to other index streams. I have talked with Steve Elliott and Mike Bethers about concentrating our efforts on six representative streams in the Haines area. I recommend the following systems:

Chilkat river (20-22 mile) 115-32-10250 Spring Creek 115-32-10250-2019-3002-4007-5002 31-mile Creek 115-32-10250-2023-3010 Kelsall river 115-32-10250-2045 Tahini river 115-32-10250-2057 Clear Creek 115-32-10280

I recently wrote a letter to Steve describing this in more detail. If you need more information please check with him. Ray Staska also mentioned that we should include the Takhin river 115-32-10250-2007. I have not surveyed this stream but he indicated it has a large number of spawning coho. Thank you for the opportunity to provide input on this subject.

State of Alaska

TO: Steve Elliott
Fisheries Biologist
Sport Fish Division
Jungan

Francy Ericksen
Francy Ericksen
Sport Fish Division
Haines

DATE: 4-1-87

FILE NO:

PECELLEGIO Salmon index streams.

throughout length.

APR 03 1987

SPORT FISH DIV.

Sorry I didn't get back to you earlier but I jist got off SLW. O.P. I designated six coho streams last year to be used as indices. Those are Chilket River (20-22mile), 31mile, Kelsall River, Spring Creek, Tahini River, and Clear Creek, I recommended dropping Nataga, Bear, and Herman Creeks from survey plans due to lack of adequate numbers of spauning who. Otherwise here goes....

| 616 , 01 |) | | | |
|------------------------|----------------------------------------------------------------------|----------------------------|-------------|--------------------------------------------------------------------------------|
| River Chilkat River | Survey Arca 20-22 mile | <u>Date</u> 11f15-12/15 | | Notes - large nos. of coho school in this area in late fall when river clears |
| Chilket River | above Tahini R, to the gorge | 10/1 - 10/30 | boat/aerial | - usually done in conjunction with |
| 31 mile | from confluence w/Klehini to incubation boxes just past 31 mile post | io/1 - 10/30 | fost | Tahin. Survey - three branches of this creek are walked at that time. |
| 37 mile | from confluence w/Klehini to 39m | | foot | - very brushy. Spawner scattered |

continued

| River | Survey Area | Date | Method | Notes |
|--------------|--------------------------------------------------------------------------------|------------------|-------------|----------------------------------------------------------------|
| Kelsall | bridge downstream two miles | 10/15-11/15 | | glacial System needs cool weather to survey |
| Little Salmm | from confluence w/ Tsirku River I mile upstream from Church Camp | 10 1 - 10 30 | flout/foot | spawners concentrate in deep wreter new camp in mid Oct. |
| Spring Creek | from confluence w/ Little Salmon upstream post bridge and major fork. | 10/1 -10/30 | fort | good chear stream |
| Tahini River | from confluence w/ Chilkat to border | 10/1 -10/30 | boat foot | glacial System but clears early. Susceptable to flooding |
| | from mouth up two miles | 10/15-11/1s | - boat/foot | good late run |

State of Alaska

Steve Elliott TO:

Fishery Biologist Division of Sport Fish Juneau

DATE:

FILE NO:

747-5355 TELEPHONE NO:

FROM:

Art Schmidt Fishery Biologist Division of Sport Fish Department of Fish and Game Sitka

Coho Index Stream SUBJECT:

April 6, 1987


Surveys


W/ MAPS (sither


Here are maps with areas surveyed for each coho index stream in the Sitka Area. These dates are dates of peak counts. Surveys prior to or after listed dates give lower numbers. Obviously you have to carefully choose the day to do the Black River helicopter flight, but it works quite well. I think Black River is the only system we could reliably count by helicopter. These are 1 inch = 1 mile, if you need to know the area of survey.


Salmon Lake has a weir so we only count the coho in the outlet below the weir just prior to removing the weir (usually mid-October).

Sorry this is so late.










DRADO 80225. OR RESTON. VIRGINIA 22092 35 AND SYMBOLS IS AVAILABLE ON PROJECT

GEOLOGICAL SURVEY

SITKA

ased on Alaska coordinate system, zone 1 Sal Transverse Mercako gost sek, dise

se Mercator projection, 1927 North American dati





To: Steve Elliott Date: August 14, 1987

Fishery Biologist
Sport Fish Division File: COSUV87

Douglas

From: Bob Zorich 33 Subject: Sport Fish Coho

Area Biologist Surveys, Petersburg Sport Fish / FRED

Petersburg

Below is a listing of areas and times I plan to conduct coho surveys for Sport Fish Division in 1987. This schedule is tentative depending on weather conditions and conflicts with coho spawning operations at Crystal Lake Hatchery.

| Stream Name ADFG Number | Area to Be Surveyed | Time | 1/ | Access |
|--------------------------------|----------------------------------------------------------------------------------------------------------------------------------|--------------|--------------|--------------------------------------|
| Bear Cr. 108-50-10030 | From Three Lakes Loop Road (TLLR) bridge to tide water. | | Q1,Q3 Q1 | One person: road access |
| Crystal Cr. 106-44-10310 | Hatchery rack returns. | | | |
| Falls Creek 106-44-10060 | From TLLR bridge to Mitkof Highway bridge. If second person is available, a survey from TLLR bridge up stream will be conducted. | Oct. Nov. | Q1, Q3 Q1 | One or Two people: road access |
| Ohmer Cr. 108-40-10500 | From upper inter-tidal to one mile above right fork and one mile above left fork. | | Q1, Q3 Q1 | One person: road access |
| Petersburg Cr. 106-44-10600 | From Petersburg Lake to upper inter-tidal. | Bet. Nov. | Q2, Q4 Q2 | One or two people: plane and boat |
| Sumner Creek: | From bridge to barrier one mile up stream. Will survey this stream same day Ohmer Creek, time permitting. | Oct. Nov. | Q1, Q3 Q1 | One person: road access |

1/ Legend: Q1- first quarter of month, Q2- second quarter of month Q3- third quarter of month, Q4- fourth quarter of month

I will be working with the Commercial Fisheries staff in Petersburg to coordinate these surveys with their's. They plan to assist me when possible and I will assist them on their surveys when possible. I will send you a list of the Commercial Fish survey plans as soon as they are available.

xc: Larson, R. Sanders Sele To: Leon Shaul

Commercial Fisheries

Douglas

From: Robert Larson /

Commercial Fisheries

Petersburg

Date: September 4, 1987

Subject: Coho Streams

The following is a list of streams in the Petersburg/Wrangell management area we are planning on surveying for coho escapements this fall.

Name and Number

Methods

Bear Creek // 108-50-03

Also on the sport fish list of target systems. Normally walked from the Three Lakes Loop Road bridge to the forks, up Canyon Creek to the falls and back to the road; the mainstem to saltwater may also be included depending upon conditions. This is a two person/two vehicle operation. Can be done in 1 day. Cost: no additional cost.

--1 ---

Falls Creek ✓ - 106-44-06 Also on the sport fish list of target systems. Normally walked from the Three Lakes Loop Road bridge downstream to the Mitkof Higway bridge. The area above the upper bridge has not been previously walked but coho distribution in this area should be documented. Two persons/two vehicles required for one day.

Timing: mid October peak. Cost: No additional cost.

Sumner Creek 108-40-40 Ohmer Creek 108-40-50 Located on the southeast portion of Mitkof Island. Also on the sport fish list of target systems. This end of Blind Slough has been the site of a remote release coho and chinook site with an intense commercial gillnet fishery at the mouth. The hatchery will no longer release coho at this site and did not release chinook there this spring. If there are no additional hatchery releases, these would be two small streams that should be included. A low priority as long as the area is flooded with chinook fry from naturally spawning adults or coho fry from Crystal Lake Hatchery.

Stikine River North Arm Ck. 108-40-10 Shakes Slough 108-40-13 Ketili Creek 108-40-14 Katete Creek 108-70-09

Kikahe River

108-40-16

Helicopter surveys can be done on one tank of fuel if conditions are good. Should be done 3 times during the fall; late September, early October, and late October. Sites such as Ketili Creek (Barnes Lake) will need to be evaluated this year, and decisions made to decide their suitability as index sites.

Cost: \$1800 per trip = \$5400

Goat Creek 108-40-17 Shuktusa Creek 108-40-18 Andrews Creek 108-40-20 Marten Creek 107-40-38 Harding River 107-40-49 N. Bradfield R. 107-40-52 E. Bradfield R. 107-40-53 Longer streams located in Bradfield Canal.
Helicopter survey from Petersburg surveyed once
with timing determined by Stikine River surveys.
Due to light conditions in late October, these will
be done on a separate day from the Stikine River.
One person/one day.
Timing: Dependent upon Stikine (mid-late October)

Cost: 2.5 hrs. flight time @ \$542 = \$1355.

Mosman Creek * 106-22-06 Navy Creek * 106-22-16 South Etolin Island streams. These are new to our proposed survey list. Survey possible with floatplane or helicopter transportation and streams walked. Each system has a barrier falls and could be walked in 2 or 3 hours less using the helicopter. Timing: walked previously in mid September. Cost: Fixed wing = \$400 plus 1 hr standby @ \$120 = \$520; Helicopter = 1.5 hrs. flight time @ \$542 plus 1 hr. standby @ \$125 = \$940

109-42-40 (Port Camden) 109-45-10 (Security Bay) 109-52-08 (Rowan Bay) These three streams are located on the northwest portion of Kuiu Island and can be accessed by vehicle from the Rowan Bay logging camp. The barrier falls and road routes are shown on the attached map. The first year I recommend using a helicopter for one trip to delineate the portions of the streams we want to use as our index areas and plan on walking only what we can do in two days. A forest service truck and bunkhouse is available in Rowan Bay for our use.

Cost: Fixed wing; round trip to Rowan Bay @ \$330 plus \$50 board; Helicopter = 2.5 hrs. flight time @ \$542 = \$1355.

Kake Bake 7 105-32-20

Duncan Creek ✓ 106-43-75 Kupreanof Island. New to survey list. Skiff access from Petersburg. Helicopter survey of this and adjacent streams required for 1987 for distribution.

Timing: mid September-October;

Cost: No additional costs when done by skiff. Helicopter: one time cost for 1987; 1 hr = \$542.00.

Portage Creek

Alternate survey site can be accessed by fixed wing or helicopter located in Portage Bay on the north side of Mitkof Island. Walk from falls to saltwater.

Cost: 1987 1 hr. helicopter time = \$542.00

Snake Creek 107-30-70 Alternate survey site can be accessed by skiff from Wrangell. Located on Etolin Island. It's known locally for a good coho return. Walk from the falls to tidewater. Two people for 1 day.


falls to tidewater. Two people for 1 day.

Cost: skiff access; \$80 per diem; \$70 airfare Petersburg
to Wrangell; Helicopter - additional .5 hr flight
time for Mosman and Navy Creek flight = \$275

| | Cost Summary | |
|----------------------|---------------------------------------------------|---------------------|
| | <u>1987</u> | <u>Future Years</u> |
| Petersburg Road Syst | em -0- | -0- |
| Stikine River | \$5400 | \$5400 |
| Bradfield Canal | \$1350 | \$1350 |
| Etolin Island | \$1215 | \$ 520 |
| Rowan Bay | \$1355 | \$ 380 |
| Kupreanof Island | <u>\$2199</u> | <u>\$340</u> |
| Total | \$11,524 | \$8,025 |
| Amount Available: C | Comm Fish - \$8,700 Sport Fish- <u>\$2,000</u> | ÷ |

\$10,700

cc: Steve Elliott Randy Timothy Robert Zorich





1,

MEMORANDUM

State of Alaska

TO: Jeff Kelly

DATE: 7/15/88

FILE NO:

TELEPHONE NO:

FHOM: Gordie Woods

SUBJECT: 1987 yelentet colo ascopement

Jeff.
attacked are the excepement counts for 1987. Counts we have
for coho are partial, and, as you will note, in some cases we got
no count at all. You will see why if you glance at the
prespectation record. We besievely have no idea what the
coho excepement was last year!

Dorde

WEATHER - 1987

A very wet winter, spring, and early summer left area systems full of water, and smolt survival and outmigration should have been good and on time. snowfall for the 1986-1987 winter was only 111 in., 47% below average. During the nine month period, October, 1986, through June, 1987, every month was above average in precipitation. Rainfall of 18.34 in. in June was the highest ever recorded for that month. July and August then set records for dryness for those two months, respectively. Juvenile mortality was very heavy during this time as many area streams dried up. Fry rescue operations were performed on Ophic Creek. Low water conditions also affected adult migrations. Humpback Creek water level was low enough to prevent pinks from entering for several weeks in August. The commercial opening on the Tsiu River was delayed as coho enturing the river had to hold within the fishery area because low water conditions prevented them from entering the lake. September set an all-time single month record for precipitation as 48.33 in. of rain fell during that month. Areawide flood conditions benefited coho escapement as gear efficiency was greatly reduced under those conditions. September's record was then erased in October as 40.37 in. of rain fell by the 27th of that month. Most coho surveys were flooded out in September and October, and for the rest of the year as well. Total precipitation for 1987 was 251 inches, making this the wettest year on record for Yakutat.

YAKUTAT PRECIPITATION AND SNOWFALL

JANUARY - DECEMBER 1987

| MONTH | PRECIP, (in.) | TOTAL SNOWFALL(in.) | PRECIP. DEVIAT | |
|----------|---------------|---------------------|-------------------|-------------------------------------|
| January | 22.95 | 31 | +13.56 | |
| February | 14.50 | 10.3 | -1- 4.45 | 33.7° mean temp., 6° above average. |
| March | 11.36 | 12.7 | + 1.81 | above average. |
| April | 15.90 | 9.7 | + 7.28 | |
| May | 15.07 | ${f r}$ | -1- 5.95 | • |
| June | 18.34 | Ø | -1-12.78 | Record wet June. |
| July | 1.86 | ø | - 6.4 <u>1</u> / | Record dry July. |
| August | 2.42 | ø | - 7.64 <u>2</u> / | Record dry August. |
| Septembe | 48.33 | Ø | +32.33 3/ | Record wet September |
| October | 48.81 | () | +28.69 4/ | Record wet October |
| November | 28.55 | 5.1 | +13.05 | |
| December | 23.16 | 44.0 | +10.18 | 22 in. snow in 2 days |
| | | | | |

1987 ANNUAL TOTAL 251.29

+115.8

New record wet year

TOTAL 1986-1987 SEASON SNOWFALL 111.7 in. (Nov. 1986-April 1987)

- 1/ Record dry July, very heavy fry loss.
- 2/ Record dry August, very heavy fry loss continues.
- 3/ Wettest month on record for Yakutat then record broken by October 1987.
- 4/ On October 21, record broken for October (previous was 36.45") and for any one year (previous record was 187.8" in 1965). New wettest month on record for Yakutat.

ALSEK RIVER WEEKLY CATCH DATA - 1987

| WEEK-DAT | E BOATS | KING | RED | COHO | PINK | СНИМ | TOTAL | DAYS |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|--------|---------------------------|------|-------|--------|---------|
| 24 6/1 | 3 18 | 1.6 | 407 | | | | 423 | .5 |
| 25 6/2 | | 206 | 3,578 | | | • | 3,784 | 2,0 |
| 26 6/2 | | 106 | 2,326 | | | | 2,432 | 3.0 |
| 27 7/0 | | 17 | 1,292 | | | • | 1,309 | 2.0 |
| 28 7/1 | | | 1,256 | ` | | 1 | 1,257 | 1.0 |
| 29 7/1 | | | 1,128 | | | | 1,128 | 2.0 |
| 30 7/2 | | | , | | | • | | |
| 31 8/0 | | | 388 | | | | 388 | 1.0 |
| 32 8/0 | | | 427 | | | | 427 | 1,0 |
| 33 8/1: | | | 341 | 5 | | 4 | 350 | 1.0 |
| 34 8/2 | | | 115 | 26 | | 5 | 146 | 3.0 |
| 8/2 | | | 29 | 68 | | 3 | 100 | 3.0 |
| 36 9/0 | | | 8 | 56 | | 4 | 69 | 3.0 |
| 37 9/1 | | | l | 2 | | 1 | 4 | 4.0 |
| 38 9/1 | | | 2 | 741 | | 440 | 1,183 | 4.0 |
| 39 9/2 | and the second s | | | 960 | | 752 | 1,712 | 4.0 |
| 40 10/ | | | 1 | 679 | | 712 | 1,392 | 4.0 |
| TOTAL: | 177 177 | 345 | 11,299 | 2,537 | | 1,922 | 16,102 | 38.5 |
| | The second secon | | | | | | | |
| ' | | | | | | | | |
| | | | 5 VI | AR COMPARISON | | | | |
| | | | 5 12 | ## 00:11 11K ±00 K | | | | |
| 1982 | 25R, 10coho | 532 | 27,389 | 6,534 | 6 | 357 | 34,811 | 42 |
| A Company of the Comp | 20R, 1 Leoho | 77 | 19,131 | 5,661 | 7 | 299 | 25,175 | 40 |
| | 22R, 11coho | 60 | 14,409 | 7,854 | 23 | 1,354 | 23,677 | 33 |
| | 18R, 6coho | 212 | 5,603 | 5,674 | . 8 | 423 | 11,920 | 33 |
| | 26R, 11coho | 476 | 24,164 | 1,331 | 13 | 537 | 26,521 | 34 |
| | 27R, 11cono 27R, 9coho | 345 | 11,299 | 2,537 | 0 | 1,922 | 16,102 | 38.5 |
| 190/ | Z/K, 9COHO | 247 | 11,277 | 4,331 | U | 1,722 | 10,102 | ر , ن د |

| DATE | AREA | KING RE | р соно | REMARKS |
|------|----------------|---------|--------|---------|
| 7/23 | Tanis #1 | 10 | 0 | Aerial |
| 7/23 | Tanis #2 | | 0 | Aerial |
| 8/10 | Tanis #1 | 1,20 | 0 | Aerial |
| 8/10 | Tanis #2 | 40 | 0 | Aerial |
| 8/14 | Basin Creek | 35 | 0 | Aerial |
| 8/14 | Cabin Creek | 22 | 0 | Aerial |
| 8/14 | Gines Creek | (| 0 | Aerial |
| 8/14 | Emile Creek | (| 0 | Aerial |
| 8/14 | Williams Creek | (|) | Aerial |

ALSEK RIVER ESCAPEMENT

| DATE | AREA | KING | RED | СОНО | REMARKS |
|-------|----------------|-------|--------|------|------------------------------------|
| 10/9 | Cabin Creek | | | 0 | Aerial |
| 10/9 | Gines Creek | | | 0 | Aerial |
| 10/9 | Emile Creek | | | 100 | Aerial |
| 10/9 | Williams Creek | | | 0 | Aerial |
| 10/29 | Klukshu Weir | 2,616 | 10,504 | | Final count, weir removed on 10/16 |

EAST RIVER WEEKLY GATCH DATA - 1987

| WEEK-DA | ATE BOATS | KING | RED | COHO | PINK | CHUM | TOTAL | PAYS |
|-------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|----------------------------------------|---------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------|
| 25 6/ 26 6/ 27 7/ 28 7/ 29 7/ 30 7/ 31 8/ 32 8/ 33 8/ 34 8/ 35 8/ 36 9/ 37 9/ 38 9/ 39 9/ | 713 8 720 7 727 9 704 14 711 27 718 31 725 60 701 74 708 89 715 59 722 51 729 21 705 16 712 10 719 11 | 2 14 7 1 15 16 13 9 15 3 | 122 77 547 626 4,313 7,911 19,600 32,580 37,566 15,220 10,225 4,218 604 50 39 23 2 | 1 1 13 24 33 319 1,340 883 343 867 968 341 | 25 47 26 15 | 3 1 1 5 14 43 160 1,354 3,269 3,034 621 1,421 546 53 | 124 91 554 630 4,330 7,929 19,644 32,663 37,674 15,431 11,901 8,827 4,521 1,014 2,327 1,537 396 | 3.0 2.0 2.0 2.0 3.0 4.0 3.0 3.0 4.0 4.0 4.0 |
| TOTAL: | 494 | 98 | 133,723 | 5,134 | 113 | 10,525 | 149,593 | 50.2 |
| 1982 1983 1984 1985 1986 1987 | 40R, 12eoho 52R, 14eoho 48R, 15eoho 66R, 19eoho 78R, 28eoho 89R, 16eoho | 81 30 22 67 109 98 | 5 YEA 97,785 82,204 39,023 185,851 76,355 133,723 | 2,026 4,891 10,875 8,148 2,769 5,134 | 428 273 851 801 332 113 | 4,668 9,566 22,419 10,576 14,285 10,525 | 104,988 96,964 73,190 206,143 93,850 149,593 | 40 29.5 27.5 42 28 50.2 |
| | | | | SCAPEMENT | T) T \ 1/2 | CHUM | REMARK | c |
| | DATE 6/04 6/07 7/03 7/08 7/12 7/15 7/19 7/23 7/26 8/02 8/10 8/14 8/17 | KTNG 33 | 1,500 1,600 1,300 1,700 6,300 5,500 5,500 8,000 10,300 17,300 24,000 25,000 29,500 34,000 | COHO | PINK | 600 | Aerial | _ |

EAST RIVER ESCAPEMENT - 1987 (continued)

| 10/9 East River 25,000 Ac | MARKS |
|---------------------------|----------------------|
| 10/9 Doame River 1,300 Ac | rial rial rial |

AKWE RIVER WEEKLY CATCH DATA - 1987

| | E BOATS | KING | RED | СОНО | PINK | CHUM | TOTAL | DAYS |
|--------------------------------|-------------------|------|-----------------------|------------|------------------------|------|-------------|-------|
| 26 6/27 | 5 | 145 | 209 | | | | 354 | 1.5 |
| 27 7/04 | | 24 | 1,692 | | | 3 | 1,719 | 1.5 |
| 28 7/11 | | 7 | 3,660 | | 1 | ĩ | 3,669 | 1.5 |
| 29 7/18 | | , | 2,533 | • | 3 | 9 | 2,545 | 2.5 |
| 30 7/25 | | 7 | 2,501 | | J | 27 | 2,535 | 1.5 |
| 31 8/01 | | , | 864 | | | 34 | 898 | 1.5 |
| 32 8/08 | | 1 | 278 | | | 10 | 289 | 2,0 |
| 33 ≪8/15 | | | 157 | 5 | 3 | 36 | 201 | 2.0 |
| 34 8/22 | | 4 | 161 | 13 | 9 | 79 | 266 | 2.0 |
| 35 8/29 | | '1 | 55 | 389 | 16 | 112 | 573 | 3.0 |
| 36 9/05 | | 1. | 19 | 1,387 | 10 | 147 | 1,554 | 3.0 |
| $\frac{30}{37}$ $\frac{9}{12}$ | | 1. | 19 | 1,307 | | 147 | 1,554 | 4.0 |
| 38 9/19 | | | | 1,633 | | 37 | 1,670 | 4.0 |
| $\frac{36}{39}$ $\frac{9}{13}$ | | | ∠ ₊ | 3,684 | | 17 | 3,705 | 4.0 |
| 40 - 10/3 | | | -1 | 821 | | 1 | 822 | 4.0 |
| -10 107. | , | | | | | _ | | |
| TOTAL: | 104 | 190 | 12,133 | 7,945 | 32 | 513 | 20,813 | 38.0 |
| | # ** | | | | | | | |
| | the specific W | | 5 YEAR | COMPARISON | | | | |
| 1982 | 9 | 129 | 4,971 | 7,014 | 129 | 82 | 12,325 | 23.5 |
| 183 | . 6 | 93 | 5,687 | 5,282 | 151 | 73 | 11,286 | 30.0 |
| 984 | 8 | 143 | 17,706 | 8,837 | 1,027 | 662 | 28,375 | 23.25 |
| 1985 | 9 | 135 | 4,938 | 4,044 | 19 | 45 | 9,181 | 21.0 |
| 1986 | 15 | 337 | 9,497 | 8,635 | 41 | 99 | 18,609 | 32.5 |
| 1987 | 12 | 190 | 12,133 | 7,945 | 32 | 513 | 20,813 | 38,0 |
| - 1490 | | | · | | • | | | |
| • | | | ESCAL | PEMENT | | | | |
| | DATE | KING | RED | CO110 | REMARKS | | | |
| | 7/03 | | 0 | | | | | |
| | 7/03 7/08 | • | 300 | | | | | |
| | | | 300 | | | | | |
| | 7/12 7/19 | | 500 | | | | | |
| | 7/23 | | 500 | | | | | |
| | 7/26 | | 800 | | | | | |
| | 8/02 | | 200 | | | | | |
| | 8/10 | 160 | 100 | | Kings bel Swanson C | | outlet, red | s in |

TTALIO RIVER ESCAPEMENT - 1987

| DATE | AREA | RED | СОНО | REMARKS | | | | |
|--------|---------------|-------|-------|---------|-----|------|----|------|
| 6/04 | New Italio | 500 | | Aerial | | | | |
| 7/03 | New Italio | 200 | | Aerial | | | | |
| 7/08 | New Italio | 600 | | Aerial | | | | |
| 7/12 | New Italio | ι,500 | | Aerial | | | | |
| 7/15 | New Italio | 250 | | Aerial | | | | |
| 7/19 | New Italio | 1,700 | | Aerial | | | | |
| 7/21 | Old Italio | 25 | | Acrial | | | | |
| 7/23 | New Italio | 1,500 | | Aerial | | | | |
| 7/26 | New Italio | 1,200 | | Aerial | | | | |
| 8/02 | New Italio | 3,250 | | Aerial | 150 | Reds | in | lake |
| 8/10 | New Tralio | 6,000 | | Aerial | 400 | Reds | in | lake |
| 8/25 | Ital,Io Lake | 300 | | | | | | |
| 9/05% | New Italio | | 100 | Aerial | | | | |
| 9/11 | New Italio | | 0 | Aerial | | | | |
| 10/9 | Italio Lake | 100 | | Aerial | | | | |
| . 10/9 | New Italio | | 400 | Aerial | | | | |
| 10/9 | Middle Italio | | 1,000 | Aerial | | | | |
| 10/9 | Old Italio | | 240 | Aerial | | | | |

SITUK - AHRNKLIN ESCAPEMENT, SURVEYS (NON-HAIN STEM SITUK RIVER)

| DATE | AREA | RED | соно | PINK | REMARKS |
|----------|-----------------|--------|-------|------|---------------|
| 7/12 | Sockeye Creek | 350 | | | Foot, at FH10 |
| 7/12 | Antilen | 150 | | | Foot, at FH10 |
| 7/15 | O.I.d. Silituk | 0 | | | Foot |
| 7/16 | West Fork | 50 | | | Foot |
| 7/17 | Autlen | 6,100 | | | Foot, boat |
| 7/18 | Autlen | 74 | | | Foot, at FH10 |
| 7/18 | Sockeye Creek | 600 | | | Foot, at FH10 |
| 7/21 | Mountain Stream | 2,000 | | | Aerial |
| 7/21 | Mountain Lake | 10,000 | | | Aerial |
| 7/30 | Sockeye Creek | 8,450 | • | | Foot |
| 8/08 | Mountain Lake | 4,000 | | | Boat |
| - 8/08 - | Mountain Stream | 2,400 | | | Foot |
| 8/09 | West Fork | 1.20 | | | Foot |
| 8/15 | West Fork | 278 | | | Foot |
| 9/02 | old Situk | 1,910 | | 800 | Foot |
| 10/7 | old Situl: | 12 | 1,010 | | Boat, 12 chum |
| 10/9 | Ant len | | 300 | | Aerial |

SITUK RIVER FLOAT COUNTS (MAIN STEM - 9 MILE BRIDGE TO SITUK LANDING)

| DATE | KENG | RED | COHO | PINK | REMARKS |
|------|-------|--------|-------|--------|-----------------------------|
| 6/09 | 3 | 35 | | | |
| 6/10 | , | 200 | | | Lower one mile |
| 6/16 | 45 | 2,040 | | | |
| 6/23 | 7 | 300 | | | Dark water, poor visibility |
| 6/27 | 266 | 3,800 | | | |
| 7/01 | 681 | 11,780 | | | |
| 7/06 | 768 | 21,130 | | | Poor visibility |
| 7/10 | 30 | 2,100 | | | Lower 2.5 miles |
| 7/14 | 1,055 | 10,110 | | | |
| 7/16 | | 3,100 | | | Lower 2.0 miles |
| 7/20 | 1,439 | 14,470 | | 3,000 | |
| 8/19 | 85 | 2,300 | 170 | 24,000 | |
| 8/27 | | 1,000 | 2,000 | | |

LOST RIVER WEEKLY CATCH DATA - 1987

| | | | | | | | | | 1 |
|--------------------------|-----------------------|----------------|------------|--------------|--------------|------------------|--------------|------------|---------|
| WEEK-DATE | BOATS | KING | RED | COHO | PINK | CHUM | TOTAL | DAYS |), |
| 25 6/20 | CLÓSED | | | | | | | | |
| 26 6/27 | CLOSED | | | | | | | | |
| 27 7/04 | CLOSED | | | | | | 07/ | י ר | ., |
| 28 7/11 | 1 | 20 | 351 | | 3 3 | | 374 814 | 2.5 3.5 | |
| 29 7/18 30 7/25 | .l. .l | 10 | 801 157 | | 38 | | 195 | 2.5 | 9 |
| 31 8/01 | , 1 | 1 | 351 | | 51 | | 403 | 2.5 | |
| 32 8/08 | Í | 2 | 86 | 14 | 2 | 1 | 105 | 3.0 | } |
| 33 8/15 | 1 | | 30 | ۷ | 3 | | 37 | 3.0 | |
| 34 - 3/ 22 | $\frac{1}{2}$ 2 | | 143 | 198 | 11 | 2 | 354 | 3.0 | 19) |
| 735 8/29 | '3 | | 48 | 234 | 2 | 13 | 297 | 3.0 | ٠., |
| 36 9/05 | 4 | | 6 | 641 | | 8 | 655 | 2.0 | |
| 37 9/12 | 4 | | 3 | 513 | | 1 4 | 517 1,107 | 4.0 | . , |
| 38 9/19 39 9/26 | $\frac{4}{3}$ | | | 1,103 747 | | 2 | 749 | 4.0 | |
| 140 10/3 | 4 | | | 192 | | 6 | 198 | 4.0 | |
| 10/// | | | | | | | | | |
| TOTAL: | 30 | 33 | 1,976 | 3,646 | 113 | 37 | 5,805 | 41.0 | .1 |
| | | | | | | | | | , / |
| | | | | | | | • | | |
| • | | | 5 YE | CAR COMPARIS | ON | | | | |
| 1982 | 5 | 10 | 5,102 | 9,955 | 700 | 13 | 15,780 | 40.0 | Ì |
| 1983 | 5 | 3 | 2,459 | 5,340 | 1,376 | 3 | 9,181 | 37.5 | 2 |
| 1984 | Z ₁ | 22 | 726 | 10,688 | 1,792 | 71 | 13,299 | 34.5 | |
| 1985 | 4 | 6 | 1,272 | 9,129 | 315 | 13 | 10,736 | 40.5 | |
| 1986 | 4 | 6 | 498 | 2,495 | 80 | 3 | 3,082 | 29.0 | 4.1 |
| 19 87 | Ľ; | 33 | 1,976 | 3,646 | 113 | 37 | 5,805 | 41.0 | - |
| | | | E | SCAPEMENT | | | | | |
| | | | | | | | | | - |
| | DATE | AREA | KING | RED | соно | REMARK | S | | ĺ |
| | 7/21 | Tawah | 2 | 0 | | Aerial | | | |
| | 7/21 | Ophir | | 200 | | Aerial | | | |
| | 9/01 | Tawah | 6 | | 236 | Boat | | | (2) |
| | 9/22 | Tawah | | | 1,500 400 | Aerial Aerial | | . * | |
| | 9/22 10/8 | Ophir Tawah | | | 5,000 | Boat, | REL Bridg | e to Op | hir |
| | TO LO | TOMOTH | | | - , | | J | • | |

YAKUTAT BAY WEEKLY CATCH DATA - 1987

| WEEK | DATE | волтѕ | KlnG | RED | СОНО | PINK | CHUM | TOTAL | DAYS |
|-----------|--------------|----------|------|--------|------------|--------|------------|------------------|--------------|
| 24 | 6/13 | 31 | 34 | 1,942 | | 1 | 6 | 1,983 | 1.0 |
| 25 | 6/20 | 52 | 87 | 2,891 | 13 | 8 | 10 | 3,009 | 1.0 |
| 26 | 6/27 | 4.1 | 31 | 2,412 | 1.1 | | 5 | 2,459 | 1.0 |
| 27 | 7/ 04 | . 33 | 48 | 1,884 | 26 | 33 | 77 | 2,068 | 1.0 |
| 28 | 7/11 | 12 | 17 | 3,120 | 82 | 77 | 12 | 3,308 | 2.5 |
| 29 | 7/18 | 2.3 | 51 | 11,071 | 298 | 1,168 | 27 | 12,615 | 3.5 |
| 30 | 7/25 | 24 | 45 | 1,438 | 454 | 217 | . 5 | 2,159 | 2.5 |
| 31 | 8/01 | 3 | 2 | 38 | 30 | 42 | 1 | 113 | 2.5 |
| 32 | 8/08 | 73 | 2 | 42 | 105 | 92 | 1 | 242 | 3.0 |
| -33 | 8/15 | . 4 | | 5 | 107 | 11 | 2 | 125 | 3.0 |
| 34 | 8/22 | ς β | 1 | 75 | 206 | 20 | 24 | 326 | 3.0 |
| 35 | 8/29 | 5 | | 23 | 183 | | 9 | 215 | 3.0 |
| 36 | 9/05 | 3 | | 2 | 220 | 1 | | 223 | 3.0 |
| 37 | 9/12 | 5 | | | 515 | 1 | 9 | 525 | 4.0 |
| 38 | 9/19 | 4 | | | 155 | | 7 | 162 | 4.0 |
| 39 | 9/26 | . 2 | ı | | 110 | | 1 | 112 | 4.0 |
| 40 | 10/3 | i | | | 5 | | | 5 | 4.0 |
| TOT∕ | M.: | 254 | 319 | 24,943 | 2,520 | 1,671 | 196 | 29,649 | 46.0 |
| | | | | | | | • | | |
| | | | | 5 YEAR | COMPARISON | | | | |
| OF STREET | | • | | | 0 710 | 0.400 | 070 | 21 022 | 17 E |
| 1982 | | ₹,3coho | 415 | 23,922 | 3,712 | 3,602 | 272 | 31,923 | 47.5 |
| 1983 | | t,4coho | 239 | 17,603 | 3,634 | 14,900 | 363 | 36,739 | 40.0 29.0 |
| 1984 | | R,5coho | 125 | 9,134 | 2,913 | 2,159 | 996 604 | 15,327 20,606 | 45.5 |
| 1981 | | R,4coho | 237 | 10,992 | 3,204 | 5,479 | 694 | | |
| 1980 | | R, Zeoho | 202 | 21,826 | 3,056 | 5,162 | 687 | 30,933 | 36.0 46.0 |
| 1987 | 521 | R,5coho | 319 | 24,943 | 2,520 | 1,671 | 196 | 29,649 | 40.0 |

| DATE | AREA | СОНО | PINK | REMARKS |
|------|-------------|------|-------|---------|
| 7/21 | Humpy Creek | | 0 | Aerial |
| 8/04 | Humpy Creek | | 650 | Boat |
| 8/10 | Humpy Creek | | 2,000 | Aerial |
| 8/31 | Humpy Creek | | 1,700 | Foot |

HARBY FISHERIES WEEKLY CATCH DATA - 1987 COMBINED CATCH

| MEEK- | -DATE | EOATS | s KINC | RED | COHO | PINK | CHUM | TOTAL | DAYS |
|----------------------------------------------|--------------------------------------------------------------|----------------------------------------------------------------|--------------------------------------|----------------------------------------------|------------------------------------|---------------------------|-------------------------------|---------------------------------------------------------|-----------------------------------------------|
| 26 27 28 29 30 31 32 | 6/27 7/04 7/11 7/18 7/25 8/01 8/08 | 3 4 3 4 3 2 2 | T.SHED | 427 925 2,812 2,420 1,318 155 | | | | 427 943 2,812 2,420 1,318 155 | 2.5 1.0 2.5 3.5 4.5 2.5 3.0 |
| 33 34 35 36 37 38 39 40 | 8/15 8/22 8/29 9/05 9/12 9/19 9/26 10/3 | NOT 2 5 7 3 HOT | FISHED FISHED FISHED FISHED | 7 2 1 | 2,163 3,153 | · . | 1 | 783 2,166 3,153 1,458 | 3.0 3.0 3.0 4.0 4.0 4.0 |
| тота | .l.: | 39 | 1 | 8,067 | 7,552 | | 1 | 15,635 | 47.5 |
| i V | | | | 5 YEA | R COMPARTSO | 4 | | | |
| 1982 1983 1984 1985 1986 | 3 20) 1 18; 5 14; 5 3 | R,7coho R,4coho R,7coho R,5coho R,6coho R,7coho | | | 5,457 18,661 16,366 3,978 | 60 139 2 33 0 | 49 13 8 12 3 1 | 30,795 13,323 23,831 23,078 8,860 15,635 | 29.0 22.0 28.0 34.5 43.5 47.5 |

| DATE | AREA | RED | COHO | REMARKS |
|------|--------------|-----|------|------------|
| 8/13 | Esker Creek | 30 | | Helicopter |
| 8/13 | Sudden River | 30 | | Helicopter |
| 8/13 | Spoon River | 150 | | Helicopter |
| 8/13 | Manby Stream | 0 | | Helicopter |

YAHTSE RIVER WEEKLY CATCH DATA - 1987

| WEE | KDATE | волтв | RED | СОНО | CHUM | TOTAL | DAYS |
|----------------------------------|------------------------------|-----------------------------|----------------|------------------------------------------------------|------------|------------------------------------------------------|----------------------------------------------|
| 34 35 | 8/22 8/29 | CLOSED 4 | 3 | 1,106 | | 1,109 | 3.0 |
| 36 37 38 39 | 9/05 9/12 9/19 9/26 | CLOSED 5 6 9 | 1 | 3,175 3,754 3,471 | 1 | 3,175 3,756 3,471 | 4.0 4.0 4.0 |
| ⊱⊸40 TO1 | 10/3 AL: | 2 26 | Z _į | 1,367 12,873 | 1 2 | 1,368 | 19.0 |
| | | | | 5 YEAR | COMPARISON | | |
| 198 198 198 198 4198 | 33 34 35 36 | 7 6 5 5 12 9 | 4 | 9,134 6,799 1,526 3,871 18,278 12,873 | 2 | 9,134 6,799 1,526 3,871 18,278 12,879 | 18.0 18.0 19.0 20.0 16.0 19.0 |

| DATE | AREA | COHO | REMARKS |
|------|----------------|-------|---------|
| 9/05 | · Yahtse | 150 | Λerial |
| 9/05 | Jetty Creek | 50 | Aerial |
| 9/05 | Pt. Riou Creek | 50 | Aerial |
| 9/11 | Yahtse | 3,000 | Aerial |
| 9/11 | Jetty Creek | 200 | Aerial |
| 9/11 | Pt. Riou Creek | 0 | Aerial |

YAKATAGA DISTRICT WEEKLY CATCH DATA TSIU -- KALIAKH RIVERS - 1987

| WEEK- | -DATE | BOATS | соно: | TSIU | , l | KALIAKH | | TOTAL | DAYS | |
|-------|-------|-------------------|-------|------------------|--------|---------|--------|--------|------------|-----|
| 34 | 8/22 | 1K | | CLOSED | | 98 | | 98 | | 2K |
| 35 | 8/29 | 37T,24K | | 12,288 | | 5,091 | 1 | 7,379 | 2T | 3K |
| 36 | 9/05 | 36T,23K | | 14,040 | | 8,332 | | 2,372 | 2T | 3K |
| 37 | 9/12 | 24T, 5K | | 3,552 | | 1,298 | | 4,850 | 2T | 4K |
| 38 | 9/19 | 28T, 10K | | 4,317 | | 890 | | 5,207 | 1T | 4K |
| 1 39 | 9/26 | 2011年1010 4年 | | 1,488 | NOT | FISHED | | 1,488 | 4 T | |
| 40° | 10/3 | 427 | 110 | OT FISHED | | FISHED | | · | | * |
| тота | la: | 429 T, 63K | | 35,685 | | 15,709 | | 51,394 | 11T | 16K |
| | W. | | 5 | YEAR COMPA | ARISON | | | | | |
| | | | | 1.E. 066 | | 15,856 | | 1,722 | 13 | |
| 1982 | | 40 | • | 45,866 19,687 | | 4,433 | | 24,120 | 10 | |
| 1983 | | 1.6 | | 50,875 | | 13,082 | | 53,957 | 21 | |
| 1984 | | 22 | | 64,601 | | 22,641 | | 37,242 | 20 | |
| 1985 | | 34 | | 19,251 | | 10,775 | | 30,026 | | 23K |
| 1986 | | 29T,27K | | 35,685 | | 15,709 | | 51,394 | | 16K |
| 1987 | | 37T,24K | | د00,00 | | 15,707 | • | , | | |
| 99°L | e. | | | | | | | | | |
| | | | | ESCAPEMEN | T | | | | | |
| | DΛTI | E AREA | | RED | COII | 0 | REMARI | ζS | | |
| | 7/0 | 4 Tsiu | | 1,000 | | | | | | |
| | 8/2: | | | | 50 | | | | markers | |
| | 8/2 | | | | 50 | | - | | markers | |
| | 8/20 | 6 Tsiu | | | 5,70 | | | | markers | |
| | 8/2 | 9 Tstu | | | 7,50 | | 4,000 | below | markers | |
| | 9/0 | | | | 8,50 | | | | | |
| | *9/2 | l Tsiu | | | 2,50 | 0 | High, | dark t | arer | |

Thight on 10/20 revealed entire one under water, no counts possible

^{*}Between 9/01 and 9/21 four attempts were made to survey the Tsiu. Flood conditions prevented any observation of fish.

To: Leon Shaul

Fisheries Biologist II

Coho Research

Douglas

From: Robert Larson Asst. Area Mgmt. Biologist

Commercial Fisheries

Petersburg

Date: December 18, 1987

Subject: '87 Coho Escapement

Surveys

The last coho survey of 1987 was completed December 1 and the completed escapement file sent to Doug Jones the next day. Copies of all the previous years' surveys for the Petersburg management area, ranked by stream and magnitude of escapement, are attached. I've also included a list of streams for our area that may be useful as index streams with significant escapement counts (surveys conducted September, October, November of more than the intertidal area). Attached also is the memo Bob Zorich, FRED/Sportfish, Petersburg, wrote concerning his survey efforts this fall. The following are my recommendations:

STIKINE RIVER

There are six tributaries in the U.S. portion of the Stikine River that have good visibility and have broad windows to make comparable escapement surveys. Kikahe River, Clear Creek, Shuktusa Branch, Shakes Creek, Andrews Creek and North Arm Creek will be good index streams, but I don't think Goat Creek has a large enough escapement to be appropriate nor is Ketili Creek appropriate due to the greatly variable escapement counts caused by timing and poor visibility. Two escapement surveys should be conducted by helicopter iwo weeks apart starting the middle of October. The cost will be \$1,000 per trip.

ETOLIN ISLAND

We have identified two streams on the southern shore of Etolin Island that will be ideal representatives of small non-lake rearing island type systems. They are Navy Creek and Flat Creek. Both these streams can be easily walked, are terminated by barrier falls, and have excellent visibility. I believe we can get total escapement counts at both sites with only one survey. The cost for fixed wing access was \$450 with the best timing early to mid October.

BRADFIELD CANAL


Coho school in the larger pools in the lower reaches of the large streams in this area and are difficult to see until mid November when the weather turns cold, the water level drops, and the fish are near their spawning sites. These streams also must be surveyed by helicopter, preferably twice, at a cost of approximately \$1500 per trip. I would continue to survey the North Fork Bradfield, Harding, and Oerns but drop the East Fork Bradfield and substitute Eagle Creek.

KUIU AND KUPREANOF ISLANDS


Streams in this area are generally difficult to survey because their small size necessitates walking them for accurate counts and the relatively large muskeg drainage areas keep the water levels from dropping rapidly after a rain keeping the water color dark. More work needs to be done in this area before streams that would make good index sites are discovered.

cc: Paul Larson Steve Elliott Bob Zorich


CLEAR CREEK COHO ESCAPEMENT




FALLS CREEK COHO ESCAPEMENT




SHAKES SLOUGH COHO ESCAPEMENT




SHUKTUSA BRANCH COHO ESCAPEMENT




NAVY CREEK COHO ESCAPEMENT









FLAT CREEK COHO ESCAPEMENT




KIKAHE CREEK COHO ESCAPEMENT



ANDREWS CREEK COHO ESCAPEMENT



BEAR CREEK COHO ESCAPEMENT



ANDREWS CREEK 108-40-20

| 69/09/10 | 1.0 | AERI AL | 0 | 0 |
|----------|------|---------|-----|-----|
| 75/09/08 | 1.0 | AERIAL | Ŏ | 0 |
| | | | - | • |
| 75/09/11 | 2.0 | AERI AL | 12 | 12 |
| 75/11/13 | L | AERI AL | 40 | 40 |
| 81/10/30 | 4.0 | FOOT | 186 | 186 |
| 82/10/21 | 3.5 | FOOT | 382 | 382 |
| 83/08/31 | WEIR | | 5 | 5 |
| 83/10/28 | L | HELO | 180 | 180 |
| 84/08/24 | WEIR | | 2 | 2 |
| 86/10/31 | L | HELO | 320 | 320 |
| 87/10/22 | L | HELO | 275 | 275 |
| 87/11/07 | L | HELO | 205 | 205 |
| 87/12/01 | L | HELO | 20 | 20 |
| | | | | |

BEAR CREEK 108-50-03

| 81/09/02 | L | FOOT | | 30 | | 30 | | |
|----------|-----|-------------|--------------|---------|---|-----|---------|------|
| 83/10/11 | 1.5 | F00T | | 63 | 4 | 67 | INDEX | AREA |
| 84/10/04 | 1.5 | F00T | | 91 | | 91 | INDEX | AREA |
| 85/10/08 | 1.7 | FOOT | | 64 | | 64 | INDEX | AREA |
| 86/10/31 | 3.5 | FOOT | | 1 | | 1 | TO UP | BRDG |
| 87/09/22 | 1.5 | FOOT | | 1 | | 1 | INDEX | AREA |
| 87/10/07 | 2.5 | FOOT | UPPER | AREA=21 | | 5 I | N INDEX | AREA |

| 87/11/07 87/12/01 | | HELO HELO | 39 30 | | 39 30 | |
|----------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------|-----------------------------------------------------------------------|-------------------------------------------------------------------|---------|-------------------------------------------------------------------------|-------------------------|
| KETILI-BARNES | <u>108-4</u> | 0-14 | | | | |
| 75/09/11 80/10/24 81/10/28 81/11/03 82/10/22 82/10/26 86/10/31 87/10/22 87/11/07 | .5 1.0 L L | FOOT | 0 567 346 124 2 7 305 45 690 85 | 3 95 | 0 567 346 124 2 10 400 45 690 95 | PARTIAL CT PARTIAL CT |
| KIKAHE 108-40- | 16 | | | | | |
| 75/09/10 75/11/13 76/11/08 77/11/18 81/10/21 81/11/07 82/10/26 83/10/28 86/10/31 87/10/22 87/11/07 | L 2.0 2.0 L L L L | AERIAL AERIAL AERIAL AERIAL AERIAL HELO HELO HELO HELO HELO HELO HELO | 0 450 50 0 80 0 201 120 290 142 129 | | 0 450 50 0 80 0 201 120 290 142 129 17 | FROZEN EXC VIS EXC VIS |
| GOAT CREEK 108 | -40-1 | Z | | | | |
| 75/09/10 80/11/11 82/10/19 84/10/18 86/09/10 86/10/31 87/10/22 | . L | AERIAL FOOT FOOT FOOT HELO HELO | 0 0 0 24 0 2 | | 0 0 0 24 0 2 | |
| SHUKTUSA BRANC | H 108 | <u>-40-18</u> | | | | |
| 81/11/07 82/09/13 82/10/19 83/09/13 83/10/15 84/10/18 86/09/10 86/10/31 87/10/22 87/11/07 87/12/01 | L 2.0 2.0 1.2 L .5 .5 L L | HELO FOOT FOOT FOOT FOOT HELO HELO HELO HELO | 0 55 135 35 75 0 0 60 115 4 | 7 | 0 55 142 35 75 0 0 70 115 4 | |

| 82/10/16 87/10/08 87/11/18 | L | AERIAL HELO HELO | 25 0 270 | 0 | 25 0 270 |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------|----------------------------------------------------------------------------|------------------------------------------------------------------------------|-----|-------------------------------------------------------------------------|
| BRADFIELD EAST | <u>FORK</u> | 107-40-53 | | | |
| 71/10/15 75/10/27 76/10/14 76/11/08 77/11/07 78/11/13 81/10/22 81/11/07 82/10/16 87/10/08 87/11/18 | 13.0 3.0 5.0 10.5 L 6.5 L | AERIAL AERIAL FOOT AERIAL | 0 3700 0 100 1500 170 0 0 15 0 | | 0 3700 0 100 1500 SPAWNING 170 0 0 15 |
| NORTH ARM 108- | 40–10 | | | | |
| 69/09/10 73/10/31 75/09/08 75/09/11 75/11/13 77/11/18 80/11/10 82/10/20 83/10/28 86/09/10 86/10/31 87/10/22 87/11/07 87/12/01 | 1.0 .5 L 1.0 .7 2.0 L L | AERIAL AERIAL AERIAL AERIAL AERIAL FOOT FOOT HELO FOOT HELO HELO HELO HELO | 0 50 0 30 75 75 12 94 30 4 95 22 23 0 | 3 1 | 0 50 0 30 75 78 13 94 30 4 95 22 23 |
| SHAKES SLOUGH | 108-4 | 0-13 | | | |
| 75/09/11 82/10/20 83/10/16 86/09/10 86/10/31 87/10/22 87/11/07 87/12/01 | .2 .7 L L L L | AERIAL FOOT FOOT FOOT HELO HELO HELO HELO | 0 72 80 0 55 7 6 | 1 | 0 73 80 0 55 7 6 |
| CLEAR CREEK 10 | 3 –40 – | 13A | | | |
| 76/11/08 82/10/20 83/10/14 84/10/18 86/10/31 87/10/22 | 1.0 1.5 L 1.3 L | AERIAL FOOT FOOT FOOT HELO HELO | 2 268 200 37 110 83 | | 2 268 200 37 110 83 |

HISIURICAL COHO ESCAPEMENTS PETERSBURG/WRANGELL AREA INDEX STREAMS

KAKE BAKE CREEK 105-32-20

| WANT DAMP AIRPRI TON SEE TA | | | | | |
|--------------------------------|------|------------------------|----------------|---------|---------------------------------|
| DATE 84/10/11 85/09/30 | | ITZ FOOT FOOT 17 | UPSTREAM 36 | DE AD 1 | 45 +24 OUT CK. 17 SAMP SEINE |
| 86/10/15 | L | FOOT | 50 | | 50 10 ABV ROAD |
| FLAT CREEK 106-22-06 | | | | | |
| 84/09/07 | 2.0 | F00T 3 | 28 | | 31 |
| 85/09/10 | | FOOT | 30 | | 30 |
| 86/09/17 | | F00T 12 | 81 | 1 | 94 |
| 87/10/09 | | FOOT | 102 | | 102 |
| NAVY CREEK 106-22-16 | | | | | |
| 73/09/17 | .3 | F00T 20 | 9 | | 29 |
| 84/09/07 | | FOOT | 9 | | 9 |
| 85/09/24 | | FOOT | 84 | 7 | 91 |
| 86/09/17 | | AERIAL | 04 | , | 0 |
| 87/10/09 | | FOOT | 57 | | 57 |
| FALLS CREEK 106-44-06 | | | | | |
| 75/10/03 | WEIR | | 74 | | 74 |
| 83/10/10 | | | 219 | 4 | 223 |
| 84/10/03 | | | 171 | • | 171 |
| 85/10/14 | | | 125 | | 125 |
| 86/10/31 | | | 1 | 4 | 5 |
| 87/09/24 | | | 21 | • | 21 |
| 87/10/30 | | | 4 | 1 | 5 |
| HARDING RIVER 107-40-49 | | | | | |
| 80/09/09 | 5.0 | FOOT | 10 | | 10 |
| 82/10/16 | | AERI AL | 50 | | 50 |
| 85/09/07 | | FOOT | 214 | 4 | 218 |
| 86/08/20 | | | 50 | | 50 |
| 87/10/08 | | HELO | 0 | | O NO SEE POOLS |
| 87/11/18 | | HELO | 70 | | 70 |
| BRADFIELD NORTH FORK 107-40-52 | | | | | |
| 71/10/15 | 1 | AFRIA | 3500 | | 3500 |
| 73/09/12 | | | 0 | | 0 |
| 75/10/27 | | | 800 | | 800 |
| 76/10/14 | | | 0 | | 0 |
| 76/11/08 | | | 2 | | 2 |
| 78/11/13 | | AERIAL | 120 | | 120 |
| 81/10/22 | | | 0 | | 0 |
| 81/11/07 | | HELO | 258 | | 258 |
| • • • • • | | | | | |

gusty strong wind kept the visibility from being good. We could see that the upper pools that held the fish during the last survey were empty. No bears or eagles observed.

108-40-20 Andrews Creek 20 live 0 dead

Excellent visibility with low water. The fish we saw were in the same area as previous surveys. They had large patches of white on their backs and appeared smaller and more siender than those observed on November 7.

108-40-18 Shuktusa Branch 0 live 0 dead

Some of the smaller beaver ponds adjacent to the creek frozen. Good visibility in the main pools and stream. No sign of fish.

108-40-16 Kikahe 17 live 0 dead

Fish distributed in the central portion of the stream. Very ragged looking with no bears and only 1 eagle in the area. Excellent visibility and low water. Approximately 6" of snow here while there is none at the river's mouth.

108-40-14 Ketili (Barnes Lake) 85 live 10 dead

Water level normal, visibility normal. Fish in pairs on riffles, old and ragged. Dead fish are bear or eagle kills. Open area of lake held 45 swans and 200 geese. No signs of fresh fish in lake or slough.

108-40-13a Clear Creek 30 live 0 dead

Excellent visibility throughout the stream. The fish were located in a side slough and two pools near the logjam. Fish very ragged looking.

108-40-13 Shakes Creek 0 live 0 dead

Similar survey conditions as survey on November 7 but no fish or signs of fish observed.

107-40-09 Glacier Creek 0 live 0 dead

No coho or sockeye observed. Upstream portion below falls good spawning but availability suspect due to 1/2 mile of rapids at the creek mouth. Poor visibility due to stream cover and poor light.

107-40-25 Oerns Creek 15 live 1 dead

Only fair visibility due to failing light despite low clear water level. Surveyed at 1:30 p.m. which is too late in the afternoon this time of year. Searched from the pool at the junction with Aarons Creek upstream to the falls. All the fish observed were in the upper riffles and pools below the rapids.

107-40-49 Harding River 70 live 0 dead

The high ridge on the southwest side of the steam prevents bright light from making the pools more visibile. Poor visibility in the lower section and normal visibility in the upper section below the rapids. With this low water level the falls/rapids area looks very passable. The area above the falls and lake was surveyed for 4 or 5 miles and no fish observed despite excellent visibility in this section. The stream was open, lots of snow and very clear water with light colored gravel substrate.

107-40-51 North Fork Bradfield 270 live 2 dead

Surveyed from the mouth to the major forks. Water was low and clear with clear skies and 6 inches of snow. Visibility was excellent on the riffles and shallow pools but the larger pools (mostly in the lower river) were too deep to see into. Fish were distributed from 3.5 miles above the mouth to 6 miles above. I don't think most of the pools in the downstream section held any fish because of the visibility and where I saw fish was associated with good spawning areas. The three largest schools of fish were near the mouth of the three tributaries in this section. The only fish I observed in the side channels or tribs were 60 in the tributary at mile 6.

107-40-53 East Fork Bradfield 160 live 0 dead

Surveyed 9 miles of stream. Terminated survey due to lack of habitat (rapids), dense fog, and no fish. The observed fish were distributed between a point 2 miles above the mouth to the bedrock canyon at mile 5. Few good spawning tributaries in this section. One seal at mile 3; little bear, wolf or eagle sign. All the fish observed were in shallow pools associated with riffies, and I don't think the deeper pools with poor visibility in the lower stream held many fish.

STIKINE RIVER TRIBUARIES
HELICOPTER SURVEY DECEMBER 1, 1987
ROBERT LARSON AND BRIAN LYNCH
COST: \$1,021.50

108-40-10 North Arm Creek 0 live 0 dead

Started the survey at the upper falls. Water clear but not low, although

108-40-18 Shuktusa 4 live 0 dead

Surveyed upstream then searched slower downstream. Total length. Four fish observed under log in highest pool below riffles. None observed at mouth or in beaverponds. Good visibility, clear, low water.

108-40-16 Kikahe 129 live 0 dead

Surveyed from the forks downstream, the same area as Will's survey, and the area I did last fall. Snowing lightly with heavy overcast, visibility good due to slow helicopter and low clear water and good snow cover. One small brown bear and three moose observed. Fish distributed throughout stream in pools. No room under banks. Three fish on shallow riffles.

108-40-14 Ketili 690 live 0 dead

No fish or signs of fish in the lake or beaver pond sloughs at the stream mouth. Fish distributed from the uppermost pool to the junction of the three forks upstream. Visibility fair with much of stream cloudy (from fish activity?) and many of the fish crowded under banks and logs. Looked fairly impressive for so small a stream.

108-40-13A Clear 39 live 0 dead

Surveyed both directions. Good visibility. Fish from just below logjam to just below plunge pool at falls. Nothing below falls or at mouth. No undercut banks.

108-40-13 Shakes 6 live 0 dead

Observed 3 fish on upstream flight under pool midway between mouth and fails. Looked more closely at pool on return trip. 6 adults, good visibility.

108-40-10 North Arm 23 live 0 dead

Poor visibility due to light snow and failing light. Fish distributed in upper half of stream with 15 present in the large pools one-quarter mile downstream from the terminal fails.

BRADFIELD CANAL NOVEMBER 18,1987
HELICOPTER SURVEY FROM PETERSBURG
COST: \$1844 FOR 3.4 HRS. BY ROBERT LARSON

107-40-08 Porterfield Ck. 0 live 0 dead

No coho or sockeye observed. Stream open, water clear and low. Good visibility despite poor light. Good spawning riffles throughout stream to just below the barrier falls.

108-40-10

The water was very clear but the extensive number of logs, stumps, and bushes prevented getting a really good view. This stream should be walked on the same day it is flown to see if it should be crossed off the helicopter survey list.

1987 COHO ESCAPEMENTS Bergmann & Minicucci Keku Strait

Heavy Creek (109-42-40) Just north of Gil Harbor 11/4/87 COST: \$45.00 Seat fare with Alaska Island Air; Kake to Petersburg (named for the logger/pilot who had a camp near here and crashed into trees just short of the water and died)

This stream is mostly bedrock with very little gravel in the first threequarters of a mile. It would be fairly easy to walk with felt-soled boots. It is a dark, muskeg drainage and should only be walked after at least three days with no rain. The stream was at about normal flow, but water had been very high within the past few days. About 300 yards upstream on the righthand side in the high flow channel of the creek is a small spring. It has a slightly sulphur smell, and I would guess it is flowing at about 1/10 cfs. It was noticeably warmer than the stream, probably between 50 and 60 degrees farenheit. There were no fry in the area. There was also one more spring about one mile upstream on the right side but it was just a trickle. About 3/4 of a mile upstream there was a sheer, 5-foot falls that was a barrier to pink salmon. About 40 pink jawbones and a chum jawbone were observed below the barrier but none were above the barrier. About 1 mile upstream there is the first major fork. It enters from the left side and is steep. A hundred yards or so above that is a large beaver dam on the right side of the stream. In the first large pool above the beaver dam a fish was observed, and it was probably a coho. We walked for another couple of hundred yards and then turned around due to time constraints. The further up the stream we got, the shallower the gradient became and more spawning and rearing habitat occured. We walked down the creek to the beaver dam and then walked through the woods on the south side of the stream to the beach. It is easier walking, mostly game trails and a few small muskegs, than the stream. We walked upstream for 1 hour and fifteen minutes and it took 40 minutes to get back to the beach. There was a doe and a buck on the beach when we came in, and we spotted another deer while walking out. About a hundred mailards were at the mouth and in the bay. Crab gear at the mouth had ADF&G #38613 on the buoys.

COHO ESCAPEMENT SURVEYS- STIKINE RIVER TRIBUTARIES
Helicopter Survey by Robert Larson
November 7, 1987 COST: \$1030.75 Temsco

108-40-20 Andrews Creek 205 live 0 dead

Surveyed downstream from the rapids below the uppermost patch of trees (same location as Will). Weather high overcast, two inches of snow on the ground, and good light. Water clear and low. Fish red with some showing white spots on tail. Only one pair on riffles, the rest scattered in smaller pools throughout the stream though fewer near the top and bottom. The largest, deepest pool contained no fish. Good survey.

108-40-14 Ketili Creek: 45 coho

All the fish were in the slough area just above the beaver dams. When the helicopter spooked them, they churned up a considerable amount of mud and made the visibility zero. No fish activity was observed in the tiny lake on the west side of the stream. A couple of eagles were present, and Barnes Lake had 8 swans and lots of ducks and geese.

108-40-16 Kikahe River: 142 coho, 25 trout

The survey was made about a mile and a half past the main fork, about 4 miles upstream on the topo map. The survey should probably be ended at the forks since it is mostly rapids above there. The survey was flown downstream with the sun at our backs. The first fish were observed about a half mile below those forks, and the last fish was seen about a mile above Red Slough. Visibility was excellent, and about 60 fish were in one school. This stream could be surveyed with a plane if the timing coincided with the fish being pooled up.

108-40-17 Goat Creek: 1 coho

Visibility was poor with partly glacial water. The one coho that was observed was about a half mile upstream.

108-40-13A Clear Creek: 83 coho (large str W of hot spring

Visiblity was excellent even in the deep pools. None of the fish were observed above the first fall which appears to be the upper limits of king salmon migration. However, from earlier surveys I have done on the grounds, the falls does not appear to be a barrier to good jumpers like coho. The survey was terminated about three quarters of a mile above the first falls. This stream could also be done from a plane with little loss in accuracy if timing was perfect.

108-40-18 Shuktusa Branch 115 coho

Visibility was excellent in the area above the rapids and poor below. All the fish observed were in two schools in the beaver ponds just above the rapids.

108-40-13 Shakes Creek 7 coho (sm cr that flows SE into Shakes Slough)

Water was clear but visibility was poor due to all the bushes. These fish were all seen in one small pool.

108-40-20 Andrews Creek: 275 coho

Visibility wqas excellent. The fish were mostly schooled in 4 large schools in the middle of the south fork. Fifteen coho were seen in the new main channel which cuts over to the other fork above the old weir site. No fish were observed in the south fork below that channel cutoff. This is another system that could be flown with a plane if conditions were right.

Harding River: 0 fish 107-40-49

North Fork of The Bradfield River: 0 fish 107-40-51

East Fork of the Bradfield River: 0 fish 107-40-53

The Harding River and both Forks of the Bradfield River were still partially glacial and visibilty was poor in all the pools. If fish had been present in significant numbers, I am sure I would have seen them since the helicopter spooks salmon so much. The surveys on the Harding and the North Fork were flown to the barriers and on the East Fork the survey was flown about 1 mile above what appears to be a barrier.

10/9/87

57 ADULTS

NAVY CREEK

106-22-16

One of the best streams found thus far for use as a coho index. Three days after heavy rain the stream is clear, normal water level, and good visibility. Very good survey; fish red but still holding in pools. fish at mouth or below ladder. Ladder in good condition. The first school 100 yards above the USGS stream gauge = 14 adults. Twenty-two adults located in three small pools midway between mouth and falls. The upper two pools at the logjam below the falls held 9 and 12 adults. No fish were present in the plunge pool below the falls. Above the first 25' barrier falls is another of equal size with no sign of fry or adults either above or below it. An old trail built by the CCC parallels the stream and facilitates the walk back to the mouth.

> Start: 9:45 a.m. End 1:15 p.m. Total time = 3.5 hrs. Robert Larson & Rexanne Elde

10/9/87

102 ADULTS

FLAT CREEK

106-22-06

Walked tidewater to barrier falls on mainstem. Observed all the fish in this stream. Easy to walk, excellent visibility. Clear and sunny. A small tributary enters from the north through a logging unit, good spawning and rearing area but too small to hold adults. Numerous wolf tracks on both Walk up from low tide to falls = 1.5 hrs, down = 1 hr. High tide makes pickup by fixed wing difficult. Recommend surveying at mid to low tide. An excellent stream for coho escapement indexing.

3 live chum. 1 live pink.

Bob Zorich

1987 COHO ESCAPEMENTS 10/22/87 WILLIAM BERGMANN **\$976** (1.8 Hours)

An aerial survey of the U.S. portion of the Stikine drainages was done on October 22. The helicopter departed Petersburg at 11:30 and returned at 1:30. The helicopter was flown directly to the border, and surveys were done downriver as index streams were encountered. It was a sunny, calm day with excellent visibility in those streams that were clear and did not have a lot of brush over the streams. The leaves had been falling, but there were still enough on the bushes and trees to impair visibility in the smaller streams. This was the second clear day and although it was cool out, it was not freezing. All of the fish observed were in pools. The following list is in survey order.

DATE

NUMBER

COMMENTS

Oct. 8 12 ADULTS PORTAGE BAY 110-16-02 COST: PART OF WILL'S BRADFILD CANAL TRIP

Very hard rain previous two weeks. Water level at falls approx. 8 feet higher than present level. Falls with 25' drop - 15'vertical; observed no fry above barrier falls. Two deep pools below falls. The plunge pool too turbulent to hold fish, the adjoining pool deep (30 yds. wide by 45 yds. No fish observed in this pool but is capable of hiding total escapement. Dropped off by helicopter in muskeg west of falls where there is a brushed trail that leads from a logging landing to the falls, a five minute walk from the muskeg. Distribution = 3 at first pool below large pool, 1 at small pool at rock bluff, 2 in pool opposite north fork, 4 adults and 4 jacks 20 yds. below north fork, 2 fish 150 yds. below fork in stump formed pool. Upper half of stream 70% bedrock ridges with coarse gravel, much dark moss, shallow pools and little spawning. The lower half of stream 80% shale-type gravel, little moss. Yellow, stained gravel. Large doilies in most pools. Creek okay to walk but would be better with a little less water and maybe later in the year to make sure there weren't fish holding in the upper pool.

Start 9:59 a.m. End 12:00 p.m. Total time = 2 hrs. 10 min. Robert Larson

1987 COHO ESCAPEMENTS 10/08/87 WILLIAM BERGMANN \$1,736 including flights to Portage Bay & Duncan

An aerial survey of some of the Bradfield Canal drainages was done on October 8. The helicopter departed Petersburg at 10:00 and returned at 12:30. It was refueled in Wrangell both going and returning. It was a sunny, calm day with normal to poor visibility in the streams because of shadows, silty water, and overstory in the smaller streams. This was the third day with no rain after extremely heavy rains. Ideally we would have waited another day or two, but the forecast called for more rain. The only fish observed were in Oerns Creek in one school. Because of the distance of the surveys, it was very frustrating to fly so far and not be able to survey more; i.e., Eagle, Anan or the Craig and Jekill. The following list is in survey order.

107-40-25 Oerns Creek: 80 coho

This stream was surveyed at 10:30 on the way south, but it had a lot of shadows and we were flying into the sun. It was resurveyed at noon on the way north with much better visibility, and 80 coho were observed in one school about 1 mile upstream.

107-40-38 Marten Creek: 0 fish

This stream was entirely in the shadows and the visibility was very poor.

To: Steve Elliot

Coho Research Biologist Sport Fish Division

Doublas

From: Bob Zorich 🏅

Area Biologist (

Petersburg

Date: November 20, 1987

File: COHOSU87

Phone: 772-3803

Subject: Cono Salmon Surveys

Petersburg Area 1987

The following is a summary of the coho salmon surveys I conducted for Sport Fish Division in the Petersburg area in 1987. Also attached are maps showing the portion of stream surveyed.

| Stream Name | ADF&G # | Date | | # Coh | ci | |
|------------------|------------|----------|-----|-------|----|------|
| Falls Creek | 106-44-006 | 9/24/87 | 21 | live | o | dead |
| | | 10/30/87 | 4 | live | 1 | dead |
| Bear Creek | 108-50-003 | 9/22/87 | 1 | live | o | dead |
| | | 10/07/87 | 26 | live | Ü | dead |
| Sumner Creek | 108-40-040 | 9/23/87 | 48 | live | o | dead |
| | | 10/22/87 | 12 | live | Ó | dead |
| Ohmer Creek | 108-40-050 | 9/23/87 | 49 | live | o | dead |
| | | 10/22/87 | 2 | live | 1 | oead |
| Petersburg Creek | 106-44-060 | 10/08/87 | 21 | live | o | dead |
| Flat Creek * | 106-22-006 | 10/09/87 | 102 | live | O | dead |

^{*} No map attached. I walked mainstream from tidewater to parrier fails.

Comments by Stream:

Falls Creek: the surveys were conducted from the upper bridge to the lower bridge to provide continuity to historical counts. This section of stream is extremely difficult to survey cono in because of the numerous, large and dark colored pools. Falls Creek is heavily stained from muskeg run off. I had wanted to survey an upper section of the stream (the first two miles above the upper bridge), but it takes at least three days of no rain for Falls Creek to drop low enough to survey. By that time it would start raining again. I would recommend that the traditional area of Falls Creek be dropped from the survey schedule and start conducting index counts above the upper bridge.

Bear Creek: This is another stream that is difficult to get a count on. Bear Creek is long and stained with muskeg colored water. The coho tend to concentrate in the pools making a count near impossible. Don Cornelius from Habitat assisted me on both surveys and on October 7, Don walked the traditional survey, while I walked an upper section. Don saw 5 coho compared to my 21 fish. I had wanted to walk the upper section a second time, but could not fit it in. Bear Creek is another stream that needs three days of no rain prior to surveying and I question the value

of the counts.

Summer Creek: Summer Creek has a large drainage but a major barrier exists on the mainstem approximately one mile above tide water. It takes three days after a rain befor the stream is low enough to walk, but the water is much clearer than most streams on the island. The left fork intercepts the mainstem about 1/4 mile below the barrier. The left fork is a highly braided unstable stream that forks again about 1/4 mile up. Most of the coho were in mainstem pools associated with large woody debri. This can be a good index stream. I recommend the survey area include the upper intertidal area (1/4 mile below the road bridge) to where the left fork enters the mainstem. Peak counts will probably be make from the last week in September through the first week in October. Survey at mid to low tide stage. The mainstem from the confluence of the left fork to the falls has very little holding or spawning habitat and is not worth surveying.

Ohmer Creek: Because Ohmer Creek has two long forks it is impractical to survey the entire system. A good index count can be made mainly in the mid to upper intertidal area, from one mile below the Ohmer Creek Campground, to one mile up each fork. Survey at the mid to low tide stage. This is pasically the traditional survey area. Trail and road systems provide good access to these areas. Peaks counts will probably occur the last of September to early October. Ohmer Creek and Sumner Creek should be walked the same day.

Petersburg Creek: It is my understanding that Petersburg Creek is no longer a coho salmon index stream. I concur with this decision.

Flat Creek: This is by far the best cono salmon index stream that I walked. The stream is easy to walk and the fish are highly visible. It is conceivable to get a total escapement count on this stream. There is a total barrier just over one mile upstream. Though most of the fish are holding in areas 1/2 mile to 1 mile up in the stream, I would recommend accessing the stream at low tide for fixed wing drop-off and pick-up. A high tide can flood the lower half mile of stream forcing you to fight the brush. Early to mid October looks to be optimum survey time. This stream can be surveyed in three hours or less.

In addition to the coho salmon stream surveys, Crystal Lake Hatchery nad a documented coho salmon escapement of 2,790 adults (1,430 males and 1,360 females) and 276 jacks.

xc: Larson

MEMORANDUM

State of Alaska

Lou Bandirola TO:

Acting Director

Division of Sport Fish

Juneau

May 1, 1987

FILE NO.:

DATE:

508.01.0

THRU:

TELEPHONE NO.:

465-4270

SUBJECT:

1986 Coho Salmon Escapements to Northern Southeast

Alaska

FROM:

Gary Sanders ##\$

Regional Research Supervisor - RI

Division of Sport Fish

Douglas

Attached are three figures illustrating the low coho salmon escapements in the northern half of southeast Alaska in 1986. Two additional figures are attached which show the effect on the Juneau marine recreational fishery for coho salmon resulting from the lack of coho salmon in the Juneau area.

The Yakutat area index streams indicated that the 1986 coho salmon escapement were the lowest in this decade (Figure 1). The 1986 coho salmon escapement (4,637) was 49.9% of the previous 6-year average of 9,287.

The six index streams in the Sitka area also received their lowest coho salmon escapements during the 1980's (Figure 2). The 1986 coho salmon escapement (31) was 28.9% of the previous 6-year average of 107. The data presented in Figure 1 also suggests a 3-year cyclic low in the coho salmon escapements in the Sitka area, i.e., 1980, 1983, and 1986.

The Juneau area index streams also received extremely poor coho salmon escapements in 1986 (Figure 3), i.e., the lowest during this decade. The 1986 coho salmon escapement (934) was 75% of the previous 6-year average of 1,239. Montana Creek, the single largest producer of coho salmon on the Juneau road system received the poorest escapement; 60 coho salmon were observed in 1986 versus the previous 6-year average of 560 coho salmon, i.e., 10.7% of the Montana Creek average coho escapement.

The poor coho salmon catch rate, catch-per-unit-effort (CPUE), in the Juneau marine recreational fishery in July 1986 (Figure 4) was a key factor influencing the 10-day closure of the commercial troll fishery in early August. Subsequently, the 1986 CPUE increased to mirror the previous 3-year mean CPUE. However, it did decline rapidly in late September. The 1986 Juneau marine recreational fishery for coho salmon ended with a below average catch (Figure 5). The 1986 sport harvest of coho salmon in the Juneau marine fishery (9,763) was 72.3% of the previous 6-year average (13,500).

As an after-thought, I have attached a table (Table 1), summarizing the data to produce Figures 1-3.

If I can supply you with additional data or clarify any of this information, just let me know.

Attachment

cc:

Frank Van Hulle




Figure 1, Coho salmon escapement data for the Yakutat area, 1930-1986.

NUMBER OF COHO




Figure 2. Coho salmon escapement data for the Sitka area, 1980-1986.

COHO ESCAPEMENT INDEX FOR SIX STREAM JUNEAU AREA 1981-1986




Figure 3. Coho salmon escapement data for the Juneau area, 1980-1986.





Figure 4. Catch per unit of effort (CPUE) data for the Juneau marine sport fishery for coho salmon.

COHO SPORT CATCH (Thousands)



JUNEAU SPORT FISHERY COHO HARVEST

Figure 5. Juneau marine recreational fishery coho salmon harvest data, 1980-1986.

Table 1. Coho salmon escapement indices in the in the Sitka, Juneau, and Yakutat areas, 1980-1986.

| YEAR | SITKA | JUNEAU | YAKUTAT |
|-------------|-------|--------|---------|
| 1980 | 50 | ND* | 6425 |
| 1981 | 116 | 1535 | 8158 |
| 1982 | 146 | 1547 | 10052 |
| 1983 | 35 | 1382 | 6383 |
| 1984 | 175 | 1361 | 10558 |
| 1985 | 122 | 1610 | 14148 |
| 1980-85 AVE | 107 | 1487 | 9287 |
| 1986 | 31 | 934 | 4637 |

*ND = No Data

MEMORANDUM

State of Alaska

DEPARTMENT OF FISH AND GAME

TO: Frank Van Hulle

Regional Supervisor Sport Fish Division

Douglas

DATE:

January 23, 1986

FILE NO .:

508.00.0

THRU:

TELEPHONE NO.:

465-4270

SUBJECT:

Complete 1985 CF/SF Coho Salmon Escapement

Data

FROM:

Gary Sanders HHA

Regional Research Supervisor

Sport Fish Division

Douglas

The following table summarizes the 1985 coho salmon escapement data collected by the Divisions of Commercial Fisheries (CF) and Sport Fish (SF):

| | ESCAPEMEN | NT | |
|-----------------------------------------------------------------|-----------|-------|-----------------------|
| AREA | COUNT | DIV. | COMMENT |
| A. Yakutat | | | |
| | | | |
| 1. Akwe River | 2,400 | CF/SF | 41% of 10-yr avg. |
| 2. Alsek River | - | | |
| a. Tannis River | 400 | CF | |
| b. Cabin Creek | 50 | CF | |
| 3. Ankau River | 5,000 | CF/SF | |
| 4. Doame River | 5,000 | CF | |
| 5. Italio River | 3,000 | CF/SF | average |
| 6. Old Italio River | 2,500 | CF | |
| 7. Tawah Creek | 15,000 | CF/SF | 324% of $7-yr$ avg. |
| 8. Lost River | 6,000 | CF/SF | 194% of 10-yr avg. |
| 9. Situk River | 6,490 | CF/SF | average |
| 10. Mamby Stream | 9,100 | CF | |
| 11. Spoon Creek | 500 | CF | |
| 12. Sudden Stream | 1,950 | CF | |
| 13. Esker Stream | 1,450 | CF | |
| 14. Jetty | 5,100 | CF | |
| 15. Kaliakh River | 37,500 | CF | |
| <pre>16. Tsiu-Tsivat Rivers</pre> | 52,350 | CF/SF | 181% of 10 -yr avg. |
| 17. Kiklukh River | 14,000 | CF | |
| B. Haines | | | |
| | | | |
| Chilkoot Lake (weir) Chilkat River | 2,188 | CF | 180% of average |
| a. Lower Chilkat River (below Wells Bridge) | 1,500 | CF | |
| b. Upper Chilkat River (above Tahini) | 41 | CF | |

| C. Juneau 1. Auke Creek (weir) 942 CF 150% of 5-yr avg. 2. Hasselborg River 550 CF 3. Jims Creek 156 CF 4. Chaik Bay Creek 104 CF 5. Switzer Creek 122 SF average 6. Salmon Creek 1,300 SF hatchery return 7. Peterson Creek (25 mi.) 276 SF above average 8. Steep Creek 186 SF average 9. Jordan Creek 70 SF poor 10. Montana/McGinnis Creeks 810 SF excellent 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 150 SF average 14. Taku River a. Sockeye Creek 150 SF average c. Yehring Creek 560 SF below average d. Fish Creek 180 SF good 2. St. John Creek 180 SF good 2. St. John Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF average 8. Kizuchia Creek 193 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | ESCAPEME | ENT | | |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|----------------------------------------|----------|--------|-------------------|--|
| d. Kelsall River | | AREA | COUNT | DIV. | COMMENT | |
| d. Kelsall River e. Tsirku River f. Spring Creek g. 37-Mile Creek g. 37-Mile Creek 65 | | · - · · - | 268 | CF | past peak | |
| f. Spring Creek g. 37-Mile Creek 6. 65 CF 3. Berners River 6,169 CF 113% of 5-yr av C. Juneau 1. Auke Creek (weir) 2. Hasselborg River 3. Jims Creek 4. Chaik Bay Creek 5. Switzer Creek 6. Salmon Creek 1.300 SF 7. Peterson Creek (25 mi.) 276 SF 8. Steep Creek 186 SF 9. Jordan Creek 186 SF 9. Jordan Creek 11. Peterson Creek (0uter Pt.) 11. Peterson Creek (Douglas Is.) 12. Fish Creek (Douglas Is.) 13. Johnson Creek 14. Taku River 15. Sockeye Creek 18. Seep 16. Sittakanay Creek 18. Seep 17. Seep 18. Sittakanay Creek 18. Seep 18. S | | · - | 132 | CF | · • | |
| g. 37-Mile Creek 65 CF 3. Berners River 6,169 CF 113% of 5-yr av. C. Juneau 1. Auke Creek (weir) 942 CF 150% of 5-yr av. 2. Hasselborg River 550 CF 3. Jims Creek 156 CF 4. Chaik Bay Creek 104 CF 5. Switzer Creek 122 SF average 6. Salmon Creek 1,300 SF hatchery return above average 8. Steep Creek 186 SF average 9. Jordan Creek 70 SF poor 10. Montana/McGinnis Creeks 810 SF excellent 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 150 SF average 14. Taku River a. Sockeye Creek 180 SF excellent 150 SF average 4. Fish Creek 180 SF good 6. Fish Creek 180 SF good 7. Sittka 1. Sinitsin Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - wild stock 86 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | ·— · | 350 | CF | | |
| 3. Berners River 6,169 CF 113% of 5-yr av. 2. Juneau 1. Auke Creek (weir) 942 CF 150% of 5-yr av. 2. Hasselborg River 550 CF 3. Jims Creek 156 CF 4. Chaik Bay Creek 104 CF 5. Switzer Creek 122 SF average 6. Salmon Creek 1,300 SF hatchery return above average 8. Steep Creek 186 SF average 9. Jordan Creek 70 SF poor 10. Montana/McGinnis Creeks 810 SF excellent 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 560 SF below average c. Yehring Creek 560 SF below average d. Fish Creek 180 SF good e. Flannigan Slough 2,320 SF excellent f. Sittakanay Creek 500 SF ? Sitka 1. Sinitsin Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | 79 | CF | | |
| C. Juneau 1. Auke Creek (weir) 942 CF 150% of 5-yr avg 2. Hasselborg River 550 CF 3. Jims Creek 156 CF 4. Chaik Bay Creek 104 CF 5. Switzer Creek 122 SF average 6. Salmon Creek 1,300 SF hatchery return 7. Peterson Creek (25 mi.) 276 SF above average 8. Steep Creek 186 SF average 9. Jordan Creek 70 SF poor 10. Montana/McGinnis Creeks 810 SF excellent 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 150 SF average 14. Taku River a. Sockeye Creek 180 SF good c. Yehring Creek 180 SF good d. Fish Creek 180 SF good d. Fish Creek 180 SF good 1. Sinitsin Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | _ | | _ | CF | | |
| 1. Auke Creek (weir) 942 CF 150% of 5-yr avg 2. Hasselborg River 550 CF 3. Jims Creek 156 CF 4. Chaik Bay Creek 104 CF 5. Switzer Creek 122 SF average 6. Salmon Creek 1,300 SF hatchery return 7. Peterson Creek (25 mi.) 276 SF above average 8. Steep Creek 186 SF average 9. Jordan Creek 70 SF poor 10. Montana/McGinnis Creeks 810 SF excellent 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 150 SF average c. Yehring Creek 180 SF good d. Fish Creek 180 SF good d. Fish Creek 180 SF good 3. Nakwasina River 408 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 192 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | 3. | Berners River | 6,169 | CF | 113% of 5-yr avg. | |
| 2. Hasselborg River 550 CF 3. Jims Creek 156 CF 4. Chaik Bay Creek 104 CF 5. Switzer Creek 122 SF average 6. Salmon Creek 1,300 SF hatchery return 7. Peterson Creek (25 mi.) 276 SF above average 8. Steep Creek 186 SF average 9. Jordan Creek 70 SF poor 10. Montana/McGinnis Creeks 810 SF excellent 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 150 SF average c. Yehring Creek 560 SF below average d. Fish Creek 180 SF good c. Yehring Creek 560 SF excellent f. Sittakanay Creek 500 SF ? Sitka 1. Sinitsin Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? |]. <u>Ju</u> | neau | | | | |
| 2. Hasselborg River 3. Jims Creek 4. Chaik Bay Creek 5. Switzer Creek 6. Salmon Creek 7. Peterson Creek (25 mi.) 8. Steep Creek 186 SF average 9. Jordan Creek 11. Peterson Creek (0uter Pt.) 11. Peterson Creek (0uter Pt.) 12. Fish Creek (Douglas Is.) 13. Johnson Creek (Douglas Is.) 13. Johnson Creek (Douglas Is.) 14. Taku River 15. Sockeye Creek 16. SF average 17. Peterson Creek (0uter Pt.) 18. Spr average 19. Johnson Creek (0uter Pt.) 19. Fish Creek (Douglas Is.) 10. Montana/McGinnis Creeks 11. Peterson Creek (Douglas Is.) 12. Fish Creek (Douglas Is.) 13. Johnson Creek (Douglas Is.) 14. Taku River 15. Sockeye Creek 15. Spr average 16. Fish Creek 15. Spr average 17. Sithanigan Slough 18. Spr good 18. Sinitsin Creek 180 SF good 180 SF good 180 CF/SF good | | | 942 | CF | 150% of 5-yr avg. | |
| 4. Chaik Bay Creek 104 CF 5. Switzer Creek 122 SF average 6. Salmon Creek 1,300 SF hatchery return 7. Peterson Creek (25 mi.) 276 SF above average 8. Steep Creek 186 SF average 9. Jordan Creek 70 SF poor 10. Montana/McGinnis Creeks 810 SF excellent 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 150 SF average c. Yehring Creek 560 SF below average d. Fish Creek 180 SF good e. Flannigan Slough 2,320 SF excellent f. Sittakanay Creek 500 SF? Sitka 1. Sinitsin Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | 550 | CF | , 0 | |
| 5. Switzer Creek 122 SF average 6. Salmon Creek 1,300 SF hatchery return 7. Peterson Creek (25 mi.) 276 SF above average 8. Steep Creek 186 SF average 9. Jordan Creek 70 SF poor 10. Montana/McGinnis Creeks 810 SF excellent 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 150 SF average c. Yehring Creek 560 SF below average d. Fish Creek 180 SF good e. Flannigan Slough 2,320 SF excellent f. Sittakanay Creek 500 SF? Sitka 1. Sinitsin Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | 156 | CF | | |
| 6. Salmon Creek 1,300 SF hatchery return 7. Peterson Creek (25 mi.) 276 SF above average 8. Steep Creek 186 SF average 9. Jordan Creek 70 SF poor 10. Montana/McGinnis Creeks 810 SF excellent 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 150 SF average c. Yehring Creek 560 SF below average d. Fish Creek 180 SF good e. Flannigan Slough 2,320 SF excellent f. Sittakanay Creek 500 SF? Sitka 1. Sinitsin Creek 109 CF/SF good 2. St. John Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | | CF | | |
| 7. Peterson Creek (25 mi.) 276 SF above average 8. Steep Creek 186 SF average 9. Jordan Creek 70 SF poor 10. Montana/McGinnis Creeks 810 SF excellent 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 150 SF average c. Yehring Creek 560 SF below average d. Fish Creek 180 SF good e. Flannigan Slough 2,320 SF excellent f. Sittakanay Creek 500 SF? Sitka 1. Sinitsin Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | 122 | SF | average | |
| 8. Steep Creek 186 SF average 9. Jordan Creek 70 SF poor 10. Montana/McGinnis Creeks 810 SF excellent 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 150 SF average c. Yehring Creek 560 SF below average d. Fish Creek 180 SF good e. Flannigan Slough 2,320 SF excellent f. Sittakanay Creek 500 SF? Sitka 1. Sinitsin Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | 1,300 | SF | hatchery return | |
| 9. Jordan Creek 70 SF poor 10. Montana/McGinnis Creeks 810 SF excellent 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 150 SF average c. Yehring Creek 560 SF below average d. Fish Creek 180 SF good e. Flannigan Slough 2,320 SF excellent f. Sittakanay Creek 500 SF ? Sitka 1. Sinitsin Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | • | SF | above average | |
| 10. Montana/McGinnis Creeks 810 SF excellent 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 150 SF average c. Yehring Creek 560 SF below average d. Fish Creek 180 SF good e. Flannigan Slough 2,320 SF excellent f. Sittakanay Creek 500 SF? Sitka 1. Sinitsin Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | 186 | SF | average | |
| 11. Peterson Creek (Outer Pt.) 144 SF excellent 12. Fish Creek (Douglas Is.) 31 SF average 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 150 SF average c. Yehring Creek 560 SF below average d. Fish Creek 180 SF good e. Flannigan Slough 2,320 SF excellent f. Sittakanay Creek 500 SF? Sitka 1. Sinitsin Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | | | poor | |
| 12. Fish Creek (Douglas Is.) 13. Johnson Creek (Douglas Is.) 14. Taku River a. Sockeye Creek b. Johnson Creek c. Yehring Creek d. Fish Creek e. Flannigan Slough f. Sittakanay Creek 150 SF good e. Flannigan Slough f. Sittakanay Creek 109 CF/SF 300 300 300 300 300 300 300 3 | | | | SF | | |
| 13. Johnson Creek (Douglas Is.) 24 SF average 14. Taku River a. Sockeye Creek 214 SF poor b. Johnson Creek 150 SF average c. Yehring Creek 560 SF below average d. Fish Creek 180 SF good e. Flannigan Slough 2,320 SF excellent f. Sittakanay Creek 500 SF? Sitka 1. Sinitsin Creek 144 CF/SF good 2. St. John Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | | SF | excellent | |
| 14. Taku River a. Sockeye Creek b. Johnson Creek c. Yehring Creek d. Fish Creek e. Flannigan Slough f. Sittakanay Creek 2,320 SF excellent f. Sittakanay Creek 100 CF/SF good CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | | SF | average | |
| b. Johnson Creek 150 SF average c. Yehring Creek 560 SF below average d. Fish Creek 180 SF good e. Flannigan Slough 2,320 SF excellent f. Sittakanay Creek 500 SF ? Sitka 1. Sinitsin Creek 144 CF/SF good 2. St. John Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | 13. | Johnson Creek (Douglas Is.) Taku River |) 24 | SF | average | |
| b. Johnson Creek c. Yehring Creek d. Fish Creek e. Flannigan Slough f. Sittakanay Creek 180 SF good e. Flannigan Slough f. Sittakanay Creek 2,320 SF excellent f. Sittakanay Creek 144 CF/SF good cF/SF good cF/SF good cF/SF good cF/SF sood cF/SF cod cF/SF cod cF/SF cod cF/SF cod cF/SF cod cF/SF cf/ | | a. Sockeye Creek | 214 | SF | poor | |
| c. Yehring Creek d. Fish Creek l80 SF good e. Flannigan Slough f. Sittakanay Creek 100 SF 2,320 SF excellent f. Sittakanay Creek 100 SF 2. Sitka 1. Sinitsin Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF yoor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 109 CF/SF yoor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - hatchery stock 109 CF/SF yoor 100 CF/SF yoor 101 January 102 CF/SF yoor 103 CF/SF yood 104 Starrigavin Creek 105 CF/SF yoor 106 Indian River - hatchery stock 107 CF/SF yood 108 Starrigavin Creek 108 CF/SF yoor 109 CF/SF yood 100 CF/SF yood 100 CF/SF yood 100 CF/SF yoor 100 SF 100 | | b. Johnson Creek | 150 | SF | • | |
| d. Fish Creek e. Flannigan Slough f. Sittakanay Creek 180 SF excellent f. Sittakanay Creek 100 SF 2. Sitka 1. Sinitsin Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF yoor 4. Salmon Lake (weir) 1. 388 SF good 5. Indian River - wild stock 6. Indian River - hatchery stock 450 CF/SF yoor 6. Indian River - hatchery stock 450 CF/SF xood 7. Starrigavin Creek 193 CF/SF xoerage 8. Kizuchia Creek 122 CF/SF yoor 188 NSERAA ? | | | 560 | SF | | |
| f. Sittakanay Creek 500 SF ? Sitka 1. Sinitsin Creek 144 CF/SF good 2. St. John Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | d. Fish Creek | 180 | SF | | |
| 1. Sinitsin Creek 144 CF/SF good 2. St. John Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | 2,320 | SF | excellent | |
| 1. Sinitsin Creek 144 CF/SF good 2. St. John Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | f. Sittakanay Creek | 500 | SF | ? | |
| 2. St. John Creek 109 CF/SF good 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | . Sit | ka | | | | |
| 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | 144 | CF/SF | good | |
| 3. Nakwasina River 408 CF/SF poor 4. Salmon Lake (weir) 1,388 SF good 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | | CF/SF | • | |
| 5. Indian River - wild stock 86 CF/SF poor 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | | | _ | |
| 6. Indian River - hatchery stock 450 CF/SF good 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | | | good | |
| 7. Starrigavin Creek 193 CF/SF average 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | | | | | poor | |
| 8. Kizuchia Creek 122 CF/SF average 9. Sea Lion Cove 188 NSERAA ? | b. | Indian River - hatchery sto | | | - | |
| 9. Sea Lion Cove 188 NSERAA ? | | | | | | |
| 10 71.4 | | | | | | |
| AVA DALLAE FOIL WAITER XX NGRDAA 7 | | | | | | |
| 11 mg 1 mg | | · | | NSERAA | ? | |
| 11. Black River 1,628 NSERAA excellent 12. Ford Arm Lake (weir) 2,325 CF average | | | | | | |

| | - | ESCAPEME | | COLOCTIVE |
|----|--------------------------------|----------|-------|-----------------------------|
| | AREA | COUNT | DIV. | COMMENT |
| Ε. | Petersburg | | | |
| | 1. Ohmer Creek | 89 | SF | poor |
| | 2. Bear Creek | 65 | SF | poor |
| | 3. Sumner Creek | 46 | SF | poor |
| | 4. Falls Creek | 125 | SF | poor |
| | 5. Petersburg Creek | 275 | SF | poor |
| 7. | <u>Ketchikan</u> | | | |
| | 1. Ward Creek - wild stock | 560 | SF | (10/24/85) below average |
| | 2. Ward Creek - hatchery stock | 1,079 | SF | (12/09/85) |
| | 3. Indian Creek | 812 | CF/SF | good |
| | 4. Carrol River | 1,550 | CF/SF | excellent |
| | 5. Eulachon River | 1,880 | CF/SF | excellent |
| | 6. Warm Chuck Lake (weir) | 956 | CF | 85% of 1982-83 avg |
| | 7. Hugh Smith Lake (weir) | 903 | CF | 54% of 1982-84 avg. |

cc: Gary Gunstrom, CF Leon Shawl, CF Art Schmidt, SF Doug Jones, CF Steve Elliott, SF